

RM# 9179

CD# 1996/048

PCT



WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification 6 :</b> C07C 257/14, C07D 213/75, 333/20, C07C 311/46, C07D 217/18, 277/40, 217/04, 231/56, 231/12, 295/135, 521/00, A61K 31/155	A1	<b>(11) International Publication Number:</b> WO 96/19440  <b>(43) International Publication Date:</b> 27 June 1996 (27.06.96)
<b>(21) International Application Number:</b> PCT/GB95/02978  <b>(22) International Filing Date:</b> 20 December 1995 (20.12.95)  <b>(30) Priority Data:</b> 9425701.1 20 December 1994 (20.12.94) GB  <b>(71) Applicant (for all designated States except US):</b> THE WELL-COME FOUNDATION LIMITED (GB/GB); Unicom House, 160 Euston Road, London NW1 2BP (GB).  <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> OPLINGER, Jeffrey, Alan [US/US]; 337 Bond Lake Drive, Cary, NC 27513 (US). GARVEY, Edward, Patrick [US/US]; 6418 Heartwood Drive, Chapel Hill, NC 27516 (US). FURFINE, Eric, Steven [US/US]; 4133 Livingstone Place, Durham, NC 27707 (US). SHEARER, Barry, George [US/US]; 203 Wintermist Drive, Cary, NC 27513 (US). COLLINS, Jon, Loren [US/US]; 106 Ledford Court, Durham, NC 27713 (US).  <b>(74) Agents:</b> CHAPMAN, Paul, William et al.; Kilburn & Strode, 30 John Street, London WC1N 2DD (GB).		<b>(81) Designated States:</b> AL, AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TT, UA, UG, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, LS, MW, SD, SZ, UG).  <b>Published</b> With international search report. With a request for rectification under Rule 91.1(f).

**(54) Title:** ACETAMIDINE DERIVATIVES AND THEIR USE AS INHIBITORS FOR THE NITRIC OXIDE SYNTHASE**(57) Abstract**

A class of acetamidine derivatives of general formula (I), methods for the manufacture thereof in therapy, particularly as inhibitors of nitric oxide synthase, is disclosed or a salt thereof, wherein  $R^1$  is hydrogen, a  $C_{1-6}$  hydrocarbyl group optionally substituted by halo, halo, nitro, cyano or a group  $XR^3$  wherein X is oxygen,  $C(O)_m$  wherein m is 1 or 2,  $S(O)_n$  wherein n is 0, 1 or 2, or a group  $NR^4$  wherein  $R^4$  is hydrogen or  $C_{1-6}$  alkyl; and  $R^3$  is hydrogen,  $C_{1-6}$  alkyl, or a group  $NR^5R^6$  wherein  $R^5$  and  $R^6$  are independently hydrogen or  $C_{1-6}$  alkyl, provided that  $R^3$  is not  $NR^5R^6$  when X is oxygen or  $S(O)_n$ ;  $R^{1a}$  and  $R^{1b}$  are independently selected from hydrogen or halo;  $R^2$  is a  $C_{1-14}$  hydrocarbyl group which may optionally contain one or two heteroatoms, the group  $R^2$  being optionally substituted by one or more groups independently selected from halo;  $N_3$ ; nitro;  $CF_3$ ;  $ZR^7$  wherein Z is oxygen,  $C(O)_m$ , wherein m' is 1 or 2,  $S(O)_n$ , wherein n' is 0, 1 or 2, or a group  $NR^8$  wherein  $R^8$  is hydrogen or  $C_{1-6}$  alkyl and  $R^7$  is hydrogen,  $C_{1-6}$  alkyl or a group  $NR^9R^{10}$  wherein  $R^9$  and  $R^{10}$  are independently hydrogen or  $C_{1-6}$  alkyl; or  $R^2$  is substituted by a group (a), wherein  $R^{11}$  has a definition the same as for  $R^1$ ; with the proviso that when  $R^1$  is a  $C_{1-6}$  alkyl group and  $R^2$  is a  $C_{1-14}$  hydrocarbyl substituted by two groups  $ZR^7$  wherein one group  $ZR^7$  is  $CO_2H$ , the other group  $ZR^7$  is not  $NH_2$ .



RTS - 0066

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

# Acetamidine derivatives and their use as inhibitors for the nitric oxide synthase

5 The present invention relates to acetamidine derivatives, to methods for their manufacture, to pharmaceutical compositions containing them and to their use in therapy, in particular their use as selective inhibitors of nitric oxide synthase.

10 It has been known since the early 1980's that the vascular relaxation brought about by acetylcholine is dependent on the presence of the endothelium and this activity was ascribed to a labile humoral factor termed endothelium-derived relaxing factor (EDRF). The activity of nitric oxide (NO) as a vasodilator has been known for well over 100 years and NO is the active component of amyl nitrite, glyceryl trinitrite and other nitrovasodilators. The recent identification of EDRF as NO has coincided with the discovery of a biochemical pathway by which NO is synthesised from the amino acid L-arginine by the enzyme NO synthase.

15 NO is the endogenous stimulator of the soluble guanylate cyclase enzyme and is involved in a number of biological actions in addition to endothelium-dependent relaxation including cytotoxicity of phagocytic cells and cell-to-cell communication in the central nervous system (see Moncada et al, Biochemical Pharmacology, 38, 1709-1715 (1989) and Moncada et al, Pharmacological Reviews, 43, 109-142 (1991)). It is now thought that excess NO production may be involved in a number of conditions, particularly conditions which involve systemic hypotension such as septic (toxic) shock and therapy with certain cytokines.

25 The synthesis of NO from L-arginine can be inhibited by the L-arginine analogue, NG-monomethyl-L-arginine (L-NMMA), and the therapeutic use of L-NMMA for the treatment of septic (toxic) shock and other types of systemic hypotension has been proposed (WO 91/04024 and GB-A-2240041). The therapeutic use of certain other NO synthase inhibitors apart from L-NMMA for the same purpose has also

been proposed in WO 91/04024 and in EP-A-0446699.

It has recently become apparent that there are at least three isoenzymes of NO synthase (reviewed in Knowles and Moncada, *Biochem. J.* (1994) 298, 249-258) as follows:

(i) a constitutive,  $\text{Ca}^{++}$ /calmodulin dependent enzyme (eNOS) which is present in vascular endothelial cells, and that releases NO in response to receptor or physical stimulation.

(ii) a constitutive,  $\text{Ca}^{++}$ /calmodulin dependent enzyme (nNOS), located in the brain and some peripheral nervous systems, that releases NO in response to receptor or physical stimulation.

(iii) a  $\text{Ca}^{++}$  independent enzyme (iNOS) which is induced after activation of vascular smooth muscle, macrophages, endothelial cells, and a number of other cells by endotoxin and cytokines. Once expressed, this inducible NO synthase synthesises NO for long periods.

The NO released by eNOS and nNOS acts as a transduction mechanism underlying several physiological responses. The NO produced by iNOS acts as a cytotoxic molecule for tumour cells and invading micro-organisms. It also appears that the adverse effects of excess NO production, in particular pathological vasodilatation and tissue damage, may result largely from the effects of NO synthesised by iNOS.

The NO synthase inhibitors proposed for therapeutic use so far, such as L-NMMA and nitroarginine, are non-selective in that they inhibit all the NO synthase isoenzymes. Use of such a non-selective NO synthase inhibitor requires that great care is taken in order to avoid the potentially serious consequences of

over-inhibition of the eNOS including hypertension and possible thrombosis and tissue damage. In particular, in the case of the therapeutic use of L-NMMA for the treatment of septic and/or toxic shock it has been recommended that the patient must be subject to continuous blood pressure monitoring throughout the treatment.


5 Thus, whilst non-selective NO synthase inhibitors have therapeutic utility provided that appropriate precautions are taken, NO synthase inhibitors which are selective in the sense that they inhibit iNOS or nNOS to a considerably greater extent than eNOS would be of even greater therapeutic benefit and much easier to use.

10 We have found that a class of acetamidine derivatives are inhibitors of NO synthase, and are useful in the treatment of systemic hypotension, and in particular the treatment of septic shock. In addition these compounds show a marked selectivity of one of the isoenzyme with little or no inhibition of the other isoenzymes.

15

Accordingly, the present invention provides an acetamidine derivative of formula (I)

20



or a salt thereof, wherein

25

$R^1$  is hydrogen, a  $C_{1-6}$  hydrocarbyl group optionally substituted by halo, halo, nitro, cyano or a group  $XR^3$  wherein X is oxygen,  $C(O)_m$  wherein m is 1 or 2,  $S(O)_n$  wherein n is 0, 1 or 2, or a group  $NR^4$  wherein  $R^4$  is hydrogen or  $C_{1-6}$  alkyl; and  $R^3$  is hydrogen,  $C_{1-6}$  alkyl, or a group  $NR^5R^6$  wherein  $R^5$  and  $R^6$  are


independently hydrogen or  $C_{1-6}$  alkyl, provided that  $R^3$  is not  $NR^5R^6$  when X is oxygen or  $S(O)_n$ ;

$R^{1a}$  and  $R^{1b}$  are independently selected from hydrogen or halo;

5

$R^2$  is a  $C_{1-14}$  hydrocarbyl group which may optionally contain one or two heteroatoms, the group  $R^2$  being optionally substituted by one or more groups independently selected from halo;  $N_3$ ; nitro;  $CF_3$ ;  $ZR^7$  wherein Z is oxygen,  $C(O)_m$  wherein  $m'$  is 1 or 2,  $S(O)_n$  wherein  $n'$  is 0, 1 or 2, or a group  $NR^8$  wherein  $R^8$  is hydrogen or  $C_{1-6}$  alkyl and  $R^7$  is hydrogen,  $C_{1-6}$  alkyl or a group  $NR^9R^{10}$  wherein  $R^9$  and  $R^{10}$  are independently hydrogen or  $C_{1-6}$  alkyl; or  $R^2$  is substituted by a group

10



15

wherein  $R^{11}$  has a definition the same as for  $R^1$ ;

20


with the proviso that when  $R^1$  is a  $C_{1-6}$  alkyl group and  $R^2$  is a  $C_{1-14}$  hydrocarbyl substituted by two groups  $ZR^7$  wherein one group  $ZR^7$  is  $CO_2H$ , the other group  $ZR^7$  is not  $NH_2$ .

25

In preferred embodiments,  $R^1$  is hydrogen, a  $C_{1-6}$  hydrocarbyl group optionally substituted by halo, halo, nitro, cyano or a group  $XR^3$  wherein X is oxygen,  $C(O)_m$  wherein m is 1 or 2,  $S(O)_n$  wherein n is 0, 1 or 2, or a group  $NR^4$  wherein  $R^4$  is hydrogen or  $C_{1-6}$  alkyl; and  $R^3$  is hydrogen,  $C_{1-6}$  alkyl, or a group  $NR^5R^6$  wherein  $R^5$  and  $R^6$  are independently hydrogen or  $C_{1-6}$  alkyl, provided that  $R^3$  is not  $NR^5R^6$  when X is oxygen or  $S(O)_n$ ; and/or

$R^{1a}$  and  $R^{1b}$  are independently selected from hydrogen or halo; and/or

$R^2$  is a  $C_{1-14}$  hydrocarbyl group which may optionally contain one or two heteroatoms, the group  $R^2$  being optionally substituted by one or two groups which may be the same or different, and are selected from halo;  $ZR^7$  wherein Z is oxygen,  $C(O)_m$  wherein  $m'$  is 1 or 2,  $S(O)_n$  wherein  $n'$  is 0, 1 or 2, or a group  $NR^8$  wherein  $R^8$  is hydrogen or  $C_{1-6}$  alkyl and  $R^7$  is hydrogen,  $C_{1-6}$  alkyl or a group  $NR^9R^{10}$  wherein  $R^9$  and  $R^{10}$  are independently hydrogen or  $C_{1-6}$  alkyl; or  $R^2$  is substituted by a group




wherein  $R^{11}$  has a definition the same as for  $R^1$ ;

with the proviso that when  $R^1$  is a  $C_{1-6}$  alkyl group and  $R^2$  is a  $C_{1-14}$  hydrocarbyl substituted by two groups  $ZR^7$  wherein one group  $ZR^7$  is  $CO_2H$ , the other group  $ZR^7$  is not  $NH_2$ .

Suitably,  $R^1$  is hydrogen,  $C_{1-4}$  alkyl, halo, nitro,  $NR^4R^3$  or  $SR^3$  wherein  $R^3$  and  $R^4$  is as hereinbefore defined.

Suitably,  $R^2$  is a  $C_{1-8}$  hydrocarbyl group optionally containing one or two heteroatoms, each group  $R^2$  being optionally substituted by one group  $ZR^7$  wherein Z and  $R^7$  are as hereinbefore defined, or a group




wherein  $R^{12}$  is fluoro, hydrogen or  $NH_2$ .


Preferably  $R^1$  is hydrogen, halo (for example fluoro) or  $NH_2$ .

Preferably  $R^{1a}$  and  $R^{1b}$  are independently hydrogen or fluoro.

5 Preferably  $R^2$  is a group




10 or




15 wherein  $p$ ,  $q$  and  $r$  are independently 0, 1 or 2 and  $R^{12}$  is as hereinbefore defined.

One embodiment of the present invention provides a compound of formula (IA)



25 wherein  $R^1$ ,  $R^{1a}$  and  $R^{1b}$  are as hereinbefore defined, and  $R^{13}$  is a  $C_{1-13}$  hydrocarbyl group which may optionally contain one or two heteroatoms, the group  $R^{13}$  being optionally substituted by one or two groups which may be the same or different, and are selected from halo,  $ZR^7$  or a group






wherein Z, R<sup>7</sup> and R<sup>11</sup> are as hereinbefore defined.


A second aspect of the present invention provides a compound of formula (IB)

5



10


wherein R<sup>1</sup>, R<sup>1a</sup> and R<sup>1b</sup> are as hereinbefore defined, and R<sup>14</sup> represents one or two substituents on the phenyl ring independently selected from halo, ZR<sup>7</sup> or a group



15

wherein Z, R<sup>7</sup> and R<sup>11</sup> are as hereinbefore defined, or R<sup>14</sup> is a C<sub>1-8</sub> hydrocarbyl group optionally containing one or two heteroatoms, and optionally substituted by one or two groups which may be the same or different, and are selected from halo, ZR<sup>7</sup> or a group

20




wherein Z, R<sup>7</sup> and R<sup>11</sup> are as hereinbefore defined.

25

In a third aspect the invention provides a compound of the formula IC:

8




5

wherein  $\text{R}^1$ ,  $\text{R}^{1a}$  and  $\text{R}^{1b}$  are as hereinbefore defined;

10  $\text{R}^{15}$  is a bond or  $\text{CH}_2$ ; and

$\text{Y}$  is  $\text{C}_{1-6}$  hydrocarbyl group, phenyl, indanyl or a 5-10 membered heterocyclic group containing 1 or 2 heteroatoms independently selected from O, N, and S;


15 wherein  $\text{Y}$  is optionally substituted by one or more groups independently selected from halo, amino, cyano, nitro, hydroxy,  $\text{C}_{1-6}$  hydrocarbyl optionally substituted by hydroxy,  $\text{C}_{1-6}$  alkoxy,  $\text{C}_{1-6}$  thioalkyl,  $\text{CF}_3$ ,  $\text{N}_3$ , a group:



20

wherein  $n$  is as hereinbefore defined,  $\text{C}(\text{O})_{m'}\text{B}$  or  $\text{S}(\text{O})_{n'}\text{B}$  wherein  $m'$  and  $n'$  are as hereinbefore defined and  $\text{B}$  is amino or  $\text{C}_{1-6}$  alkyl, a group:

25



5


wherein n is as hereinbefore defined and D and E are independently hydrogen, hydroxy, amino, C<sub>1-6</sub> alkyl, -CNHCH<sub>3</sub>, -CO<sub>2</sub>Bu, or a 5-10 membered heterocyclic group containing 1 or 2 heteroatoms independently selected from O, N or S, a group:

10



wherein n and n' are as hereinbefore defined and F is phenyl, or a 5- or 6-membered heterocyclic group containing 1 or 2 heteroatoms selected from O, N and S optionally substituted by

15



20

a group:



wherein n is as hereinbefore defined and G is -CO<sub>2</sub>H, -CO<sub>2</sub>C<sub>1-6</sub> alkyl or a 5-10 membered heterocyclic group containing 1 or 2 heteroatoms independently selected from O, N or S optionally substituted by amino, CNHNH<sub>2</sub> and CH<sub>2</sub>SCNHNH<sub>2</sub>.


25

In a preferred embodiment of this aspect of the invention a compound of the formula IC is provided wherein:

R<sup>15</sup> is a bond or CH<sub>2</sub>;

Y is C<sub>1-6</sub> hydrocarbyl, phenyl, indanyl or a 5-10 membered heterocyclic group containing 1 or 2 heteroatoms independently selected from O, N or S;


5 wherein Y is optionally substituted by one or more groups independently selected from halo, amino, cyano, nitro, hydroxy, C<sub>1-6</sub> hydrocarbyl optionally substituted by hydroxy, C<sub>1-6</sub> thioalkyl, a group:



10

wherein n is as hereinbefore defined, C(O)<sub>m</sub>.B or S(O)<sub>n</sub>.B wherein m' and n' are as hereinbefore defined and B is amino or C<sub>1-6</sub> alkyl, a group:


15



20 wherein n is as hereinbefore defined and D and E are independently hydrogen, hydroxy, amino, C<sub>1-6</sub> alkyl, -CNHCH<sub>3</sub>, -CO<sub>2</sub>'Bu, or a 5-10 membered heterocyclic group containing 1 or 2 heteroatoms independently selected from O, N or S, a group:




25 wherein n and n' are as as hereinbefore defined and F is phenyl, or a 5- or 6-membered heterocyclic group containing 1 or 2 heteroatoms selected from O, N and S optionally substituted by



5

a group:



10

wherein n is as hereinbefore defined and G is a 5-10 membered heterocyclic group containing 1 or 2 heteroatoms independently selected from O, N or S optionally substituted by amino, CNH<sub>2</sub> and CH<sub>2</sub>SCNH<sub>2</sub>.

In a particularly preferred embodiment of this aspect of the invention the invention provides a compound of the formula IC wherein:

15


R<sup>15</sup> is a bond or CH<sub>2</sub>:

Y is C<sub>1-6</sub> hydrocarbyl, phenyl, indanyl or a 5-10 membered heterocyclic group containing 1 or 2 heteroatoms independently selected from O, N or S;

20


wherein Y is optionally substituted by one or more groups independently selected from amino, C<sub>1-6</sub> hydrocarbyl optionally substituted by hydroxy, a group:

25



wherein n is as hereinbefore defined, a group:

12



5


wherein n is as hereinbefore defined and D and E are independently hydrogen, hydroxy, amino, C<sub>1-6</sub> alkyl, -CNHCH<sub>3</sub>, or a 5-10 membered heterocyclic group containing 1 or 2 heteroatoms independently selected from O, N or S, a group:

10



wherein n and n' are as hereinbefore defined and F is phenyl, or a 5- or 6-membered heterocyclic group containing 1 or 2 heteroatoms selected from O, N and S optionally substituted by

15



20

a group:



wherein n is as hereinbefore defined and G is a 5-10 membered heterocyclic group containing 1 or 2 heteroatoms independently selected from O, N or S, CNHNH<sub>2</sub> and CH<sub>2</sub>SCNHNH<sub>2</sub>.

25

A preferred group of compounds within formula (IC) are those where R<sup>15</sup> is a bond. Such compounds are selective inhibitors of the neuronal NOS. Particularly preferred compounds within this sub-group are those where Y is optionally

substituted phenyl or a optionally substituted 5-10 membered heterocyclic group containing 1 or 2 heteroatoms independently selected from O, N or S. Examples of such compounds include:

- 5     N-(3-(Hydroxymethyl)phenyl)Acetamidine;  
      N-(3-(Aminomethyl)phenyl)acetamidine;  
      N-(3-((1,2,3,4-tetrahydroisoquinol-2-yl)methyl)phenyl)-acetamidine;  
      N-(3-((Methylamino)methyl)phenyl)acetamidine;  
      N-(Dimethylamino)methyl)phenyl)acetamidine;  
10    N-(3-Morpholinomethyl)phenyl)acetamidine;  
      N-(3-(Aminomethyl)phenyl)-2-fluoroacetamidine;  
      N-(3-(2-Aminoethyl)phenyl)acetamidine; and  
      N-(1,2,3,4-Tetrahydroisoquinolin-7-yl)acetamidine;  
15    and salts thereof.

A second preferred group of compounds within formula (IC) are those where  $R^{15}$  is  $-CH_2-$ . This group of compounds are selective inhibitors of the inducible NOS. Particularly preferred compounds within this subgroup are those where Y is is optionally substituted phenyl, indanyl or a optionally substituted 5-10 membered heterocyclic group containing 1 or 2 heteroatoms independently selected from O, N or S. Examples of such compounds include:

- 20    N-(3-(aminomethyl)benzyl)acetamidine;  
25    N-(4-(aminomethyl)benzyl)acetamidine;  
      N-(3-(Aminomethyl)benzyl)-2-fluoroacetamidine;  
      N-(3-(Aminomethyl)benzyl)-2-aminoacetamidine;  
      N-[[6-(Aminomethyl)-2-pyridyl]methyl]acetamidine;  
      N-(3-(Aminomethyl)benzyl)-2,2,2-trifluoroacetamidine;

N-((1H-Indazol-6-yl)methyl)acetamidine; and  
N-((3-Amino-5-indanyl)methyl)acetamidine;

and salts thereof.

5

Especially preferred compounds within this group include:

N-(3-(aminomethyl)benzyl)acetamidine;  
N-(4-(aminomethyl)benzyl)acetamidine;  
10 N-(3-(Aminomethyl)benzyl)-2-fluoroacetamidine;  
N-(3-(Aminomethyl)benzyl)-2-aminoacetamidine;  
N-[[6-(Aminomethyl)-2-pyridyl]methyl]acetamidine;  
N-((1H-Indazol-6-yl)methyl)acetamidine; and  
N-((3-Amino-5-indanyl)methyl)acetamidine;

15

and salts thereof.

Preferred compounds of formula (I) or (IC) include:

20 N-(3-(aminomethyl)benzyl)acetamidine;  
N-(3-(aminomethyl)benzyl)-2-fluoroacetamidine;  
N-(4-(aminomethyl)benzyl)acetamidine;  
N-(4-aminobutyl)acetamidine;  
N-(5-aminopentyl)acetamidine;  
25 N-(6-aminoethyl)acetamidine;  
N-(7-aminoheptyl)acetamidine;  
N,N'-(1,3-phenylenebis (methylene))bis(acetamidine);  
N,N'-(1,4-phenylenebis (methylene))bis(acetamidine);  
N,N'-(1,3-phenylene (diethylene))bis(acetamidine);



N,N'-(1,4-phenylene (diethylene))bis(acetamidine);  
N-(3-(aminomethyl)benzyl)-2-methoxyacetamidine;  
2-Amino-N-(3-(aminomethyl)benzyl)acetamidine;  
N-(3-(hydroxymethyl)phenyl)acetamidine;  
5 N-(3-(aminomethyl)phenyl)acetamidine;  
N-(3-(Aminomethyl)phenyl)acetamidine;  
N-(3-(Aminomethyl)phenyl)-2-fluoroacetamidine; and  
N-((3-Amino-5-indanyl)methyl)acetamidine

10 and salts thereof.

Especially preferred compounds include N-(3-(aminomethyl)benzyl)acetamidine;  
N-(3-(aminomethyl)benzyl)-2-fluoroacetamidine;  
N-(3-(Aminomethyl)phenyl)acetamidine;  
15 N-(3-(Aminomethyl)phenyl)-2-fluoroacetamidine; and  
N-((3-Amino-5-indanyl)methyl)acetamidine

and salts thereof.

20 By the term "hydrocarbyl" is meant a group containing only hydrogen and carbon atoms, and may contain double and/or triple bonds and which may be straight, branched, cyclic or aromatic in nature.

The term "heteroatom" has the meaning known to one skilled in the art and  
25 includes nitrogen, oxygen and sulphur atoms.

By the term "halo" is meant fluoro, chloro, bromo, or iodo, and preferably fluoro.

The compounds of formula (I) may include a number of asymmetric centres in the

molecule depending on the precise meaning of the various groups, and formula (I) is intended to include all possible isomers.

5 As hereinbefore mentioned, the compounds of formula (I) have activity as inhibitors of NO synthase and accordingly a further aspect of the present invention provides a compound of formula (I) for use in medicine.

10 A yet further aspect provides the use of a compound of the invention in the manufacture of a medicament for the treatment of a condition requiring inhibition of NO synthase. More specifically, there is provided the use of a compound of the invention for the manufacture of a medicament for the treatment of a condition where there is an advantage in inhibiting an NO synthase isoenzyme with little or no inhibition of the other isoenzymes.

15 In another aspect, the present invention provides a method of treating a condition requiring inhibition of NO synthase, in particular the inhibition of one NO synthase isoenzyme with little or no inhibition of the other isoenzymes, comprising administering to a mammal in need thereof an effective amount of a compound of the invention or a pharmaceutically acceptable salt thereof.

20 In a further aspect, there is provided the use of a compound of the invention or salt thereof in the manufacture of a medicament for the treatment of shock states resulting from overproduction of NO by iNOS such as septic shock, or shock caused by fulminant hepatic failure or by therapy with cytokines such as TNF, IL-1 and IL-2 or therapy with cytokine-inducing agents, for example 5, 6-dimethylxanthenone acetic acid.

Other conditions where there is an advantage in selectively inhibiting iNOS include a wide range of auto-immune and/or inflammatory diseases, such as those of the

joint (e.g. rheumatoid arthritis, osteoarthritis), of the gastrointestinal tract (e.g. ulcerative colitis and other inflammatory bowel diseases, gastritis and mucosal inflammation resulting from infection), of the lung (adult respiratory distress syndrome, asthma), of the heart (myocarditis), of the nervous tissue (e.g. multiple sclerosis), of the pancreas (e.g. diabetes melitus), of the kidney (e.g. glomerulonephritis), of the skin (e.g. dermatitis, psoriasis, urticaria) as well as of transplanted organs (rejection) and multi-organ diseases (e.g. systemic lupus erythematosus). Furthermore there is evidence for overproduction of NO by iNOS in atherosclerosis. Therefore, a yet further aspect of the present invention provides the use of a compound of formula (I) or salt thereof in the manufacture of a medicament for use in treating the above conditions.

Inhibition of neuronal NOS is of benefit in the treatment of diseases of the nervous system due to over production of NO by this isoenzyme, particularly the treatment of cerebral ischemia. Other diseases include CNS trauma, epilepsy, AIDS dementia, chronic neurodegenerative disease and chronic pain, and conditions in which non-adrenergic non-cholinergic nerve may be implicated such as priapism, obesity and hyperphagia. Accordingly the present invention also provides the use of a compound of formula (I) or a salt thereof in the manufacture of a medicament for use in treating the above conditions.

Furthermore inhibition of NO synthase may be of advantage in preventing the lymphocyte loss associated with HIV infection, and in reducing tumour growth and metastasis.

Inhibition of both iNOS and nNOS may be of benefit in the treatment of certain conditions where both isoenzymes play a role. for example CNS conditions such as cerebral ischemia.

As used herein, reference to "treatment" of a patient is intended to include prophylaxis; the term "mammal" is intended to include a human or an animal.

5 The present invention includes compounds of formula (I) in the form of salts, in particular acid addition salts. Suitable salts include those formed with both organic and inorganic acids. Such acid addition salts will normally be pharmaceutically acceptable although salts of non-pharmaceutically acceptable salts may be of utility in the preparation and purification of the compound in question. Thus, preferred salts include those formed from hydrochloric, hydrobromic, sulphuric, citric, 10 tartaric, phosphoric, lactic, pyruvic, acetic, trifluoroacetic, succinic, oxalic, fumaric, maleic, oxaloacetic, methanesulphonic, ethanesulphonic, p-toluenesulphonic, benzenesulphonic and isethionic acids. Salts of compounds of formula (I) can be made by reacting the appropriate compound in the form of the free base with the appropriate acid.

15 Whilst it may be possible for the compounds of the present invention to be administered as the raw chemical, it is preferable to present them as a pharmaceutical formulation. According to a further aspect, the present invention provides a pharmaceutical formulation comprising a compound of formula (I) or 20 a pharmaceutically acceptable salt or solvate thereof, together with one or more pharmaceutically acceptable carriers therefor and optionally one or more other therapeutic ingredients, for example an antibiotic, and/or a volume replacement liquid. The carrier(s) must be "acceptable" in the sense of being compatible with the other ingredients of the formulation and not deleterious to the recipient thereof.

25 The formulations include those suitable for oral, parenteral (including subcutaneous, intradermal, intramuscular, intravenous and intraarticular), rectal and topical (including dermal, buccal, sublingual and intraocular) administration although the most suitable route may depend upon for example the condition and

disorder of the recipient. The formulations may conveniently be presented in unit dosage form and may be prepared by any of the methods well known in the art of pharmacy. All methods include the step of bringing into association a compound of formula (I) or a pharmaceutically acceptable salt or solvate thereof ("active ingredient") with the carrier which constitutes one or more accessory ingredients. In general the formulations are prepared by uniformly and intimately bringing into association the active ingredient with liquid carriers or finely divided solid carriers or both and then, if necessary shaping the product into the desired formulation.

Formulations of the present invention suitable for oral administration may be presented as discrete units such as capsules, cachets or tablets each containing a predetermined amount of the active ingredient; as a powder or granules; as a solution or a suspension in an aqueous liquid or a non-aqueous liquid; or as an oil-in-water liquid emulsion or a water-in-oil liquid emulsion. The active ingredient may also be presented as a bolus, electuary or paste.

A tablet may be made by compression or moulding, optionally with one or more accessory ingredients. Compressed tablets may be prepared by compressing in a suitable machine the active ingredient in a free-flowing form such as a powder or granules, optionally mixed with a binder, lubricant, inert diluent, lubricating, surface active or dispersing agent. Moulded tablets may be made by moulding in a suitable machine a mixture of the powdered compound moistened with an inert liquid diluent. The tablets may optionally be coated or scored and may be formulated so as to provide slow or controlled release of the active ingredient therein.

Formulations for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostats and solutes which render the formulation isotonic with the blood of the intended

recipient; and aqueous and non-aqueous sterile suspensions which may include suspending agents and thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example sealed ampoules and vials, and may be stored in a freeze-dried (lyophilised) condition requiring only the addition  
5 of the sterile liquid carrier, for example, saline, water-for-injection, immediately prior to use. Alternatively, the formulations may be presented for continuous infusion.

Extemporaneous injection solutions and suspensions may be prepared from sterile  
10 powders, granules and tablets of the kind previously described.

Formulations for rectal administration may be presented as a suppository with the usual carriers such as cocoa butter or polyethylene glycol.

15 Formulations for topical administration in the mouth, for example buccally or sublingually, include lozenges comprising the active ingredient in a flavoured basis such as sucrose and acacia or tragacanth, and pastilles comprising the active ingredient in a basis such as gelatin and glycerine or sucrose and acacia.

20 Preferred unit dosage formulations are those containing an effective dose, as herein below recited, or an appropriate fraction thereof, of the active ingredient.

It should be understood that in addition to the ingredients particularly mentioned above, the formulations of this invention may include other agents conventional in  
25 the art having regard to the type of formulation in question, for example those suitable for oral administration may include flavouring agents.


For each of the aforementioned conditions, the compounds of the invention may be administered orally or via injection at a dose of from 0.1 to 250mg/kg per day.

The dose range for adult humans is generally from 5mg to 17.5g/day, preferably 5mg to 2g/day and most preferably 10mg to 1g/day, for example 15mg to 600mg/day. Tablets or other forms of presentation provided in discrete units may conveniently contain an amount of compound of the invention which is effective at such dosage or as a multiple of the same, for instance, units containing 5mg to 500mg, usually around 10mg to 200mg.


The compounds of formula (I) are preferably administered orally or by injection (intravenous or subcutaneous). The precise amount of compound administered to a patient will be the responsibility of the attendant physician. However the dose employed will depend on a number of factors, including the age and sex of the patient, the precise disorder being treated, and its severity. Also the route of administration may vary depending on the condition and its severity.

Certain compounds of formula (I) are novel, and accordingly a further aspect of the present invention provides a process for the preparation of a compound of formula (I) as hereinbefore defined.

Thus, compounds of formula (I) or protected derivatives thereof may be prepared by the reaction of an amine of formula (II)



or a protected derivative thereof wherein  $\text{R}^2$  is as hereinbefore defined, with a 1-benzylthioethaniminium derivative of formula (III)



wherein  $R^1$ ,  $R^{1a}$  and  $R^{1b}$  are as hereinbefore defined followed by deprotection if necessary. The reaction may be carried out in a polar solvent, for example a  $C_{1-4}$  alkanol such as ethanol at a non-extreme temperature of from  $-20^\circ\text{C}$  to  $100^\circ\text{C}$  such as  $0^\circ\text{C}$ . The 1-benzylthioethaniminium derivative of formula (III) is suitably a halide derivative, preferably a chloride or bromide. Deprotection may be carried out by methods well known to those skilled in the art, for example by methods described in Greene, T. W.; Wuts, P.G.M. (1991) *Protective Groups in Organic Synthesis*, p 327 - 330, John Wiley, New York.

- 10 Compounds of formula (III) may be prepared by the S-alkylation of thioamide compounds of formula (IV)



15

wherein  $R^1$ ,  $R^{1a}$  and  $R^{1b}$  are as hereinbefore defined with benzyl bromide or chloride as described previously (Takido, Y., Itabashi, K. *Synthesis* 1987, p 817-819). The reaction may be carried out in a solvent such as chloroform at non-extreme temperatures of from  $0^\circ\text{C}$  to  $200^\circ\text{C}$ , such as refluxing chloroform.


20

Compounds of formula (II) and (IV) are commercially available or may be prepared by methods known to a person skilled in the art.

- 25 Where Y is an indanyl group, the following synthetic schemes can be used for the preparation of such compounds:




23



Reagents: (a) EtOH, H<sub>2</sub>/Pd/C; (b) NaNO<sub>2</sub>, then CuCN; (c) pyridine, BnONH<sub>2</sub>.HCl; (d) BH<sub>3</sub>.THF; (e) EtOH, 20°C; (f) (BOC)<sub>2</sub>-O, MeOH/CH<sub>2</sub>Cl<sub>2</sub>, Et<sub>3</sub>N; (g) HCl, Dioxane.

15



Reagents: (a) EtOH, H<sub>2</sub>/Pd/C; (b) pyridine, BnONH<sub>2</sub>.HCl; (c) BH<sub>3</sub>.THF; (d) (BOC)<sub>2</sub>-O, MeOH/CH<sub>2</sub>Cl<sub>2</sub>, Et<sub>3</sub>N; (e) EtOH, 20°C; (f) HBr/ AcOH.

25

An additional method for the preparation of compounds of the invention involves the conversion of an azide into an amidine utilising a one-pot method involving triphenylphosphine reduction followed by *in situ* amidination of the intermediate amine. A further refinement of the method involves the use of polymer-supported triphenylphosphine, the byproduct oxide of which can be readily filtered from the

reaction mixture.

Preferred features of each aspect of the invention are as for each other aspect, *mutatis mutandis*.

5

The present invention will now be described by way of example only:-

**Intermediate A**

10     **Preparation of S-Benzylthioacetimidate Hydrochloride** (Takido, Y.; Itabashi, K. Synthesis 1987, p 817-819)

15     A solution of thioacetamide (60.0 g) and 91.9 ml benzyl chloride in 600 ml chloroform was stirred at reflux for 14 h. Upon cooling to 0°C, the white solids were collected via filtration and dried in vacuo to yield 101 g of a white solid. Dilution of the filtrate with 400 ml Et<sub>2</sub>O and filtration provided an additional 46 g of a pale yellow solid. <sup>1</sup>H NMR (300 MHz, DMSO) δ 7.5-7.3 (m, 5H), 4.66 (s, 2H), 2.67 (s, 3H).

20     The following intermediates were prepared by an analogous method:

B     **S-Benzylthioacetimidate Hydrobromide**: from thioacetamide and benzyl bromide. <sup>1</sup>H NMR (200 MHz, DMSO) δ 2.65 (3H, s), 4.63 (2H, s), 7.51-7.31 (5H, m).

25

C     **S-(2-Naphthylmethyl)thioacetimidateHydrobromide**: from thioacetamide and 2-(bromomethyl)naphthalene. <sup>1</sup>H NMR (200 MHz, DMSO) δ 11.9 (br s, 1H), 8.04-7.92 (m, 4H), 7.6-7.55 (m, 3H), 4.8 (s, 2H), 2.68 (s, 3H).

- D S-Benzyl-2-fluorothioacetimidate Hydrobromide:** from fluorothioacetamide and benzyl bromide.  $^1\text{H}$  NMR (300 MHz, DMSO)  $\delta$  7.5-7.3 (m, 5H), 4.66 (s, 2H), 2.67 (s, 3H).
- 5 **E S-(2-Naphthylmethyl)-2-fluorothioacetimidate Hydrobromide:** from fluorothioacetamide and 2-(bromomethyl)naphthalene.  $^1\text{H}$  NMR (200 MHz, DMSO)  $\delta$  7.91-7.82 (m, 4H), 7.65-7.48 (m, 3H), 4.73 (d,  $J = 47.2$  Hz, 2H), 3.9 (s, 3H).
- 10 **F S-(2-Naphthylmethyl)-2,2-difluorothioacetimidate Hydrobromide:** from difluorothioacetamide and 2-(bromomethyl)naphthalene. (Crude)
- G S-Benzyl-2-methoxythioacetimidate Hydrobromide:** from 2-methoxy-thioacetamide and benzyl bromide.  $^1\text{H}$  NMR (200 MHz, DMSO)  $\delta$  3.42 (3H, s), 4.61 (2H, s), 4.67 (2H, s), 7.44 (5H, m).
- 15
- H S-(2-Naphthylmethyl)-2-(thiomethyl)thioacetimidate Hydrobromide:** from 2-(thiomethyl)thioacetamide and 2-(bromomethyl)naphthalene.  $^1\text{H}$  NMR (200 MHz, DMSO;  $\text{D}_2\text{O}$ )  $\delta$  2.09 (3H, s), 3.89 (2H, s), 4.67 (2H, s), 7.91-7.44 (7H, m).
- 20
- I S-Benzyl-2-tertbutylcarbamoylethioacetimidate Hydrobromide:** from 2-(tert-butylcarbamoylethio)acetamide and benzyl bromide.  $^1\text{H}$  NMR (200 MHz, DMSO)  $\delta$  1.40 (9H, s), 4.27 (2H, d), 4.56 (2H, s), 7.42 (5H, m), 7.79 (1H, t).
- 25
- J S-Benzylthiopropionimidate Hydrobromide:** from benzyl bromide and thiopropionamide.  $^1\text{H}$  NMR (400 MHz, DMSO)  $\delta$  7.46-7.32 (m, 5H), 4.59 (s, 2H), 2.83 (d,  $J = 7.5$  Hz, 2H), 1.21 (t,  $J = 7.5$  Hz, 3H).

### Example 1

#### Preparation of N-(3-(aminomethyl)benzyl)acetamide

##### Preparation of Intermediate *tert*-butyl-N-(3-(aminomethyl) benzyl)carbamate

10 g (73.42 mmol) of m-xylenediamine was added to 5.1 ml (36.71 mmol) of  
5 triethylamine and 200 ml of anhydrous methanol. To this solution at 0 °C was  
added dropwise over 60 minutes a solution of 8.0 g (36.71 mmol) of  
di-*tert*-butyldicarbonate in 60 mL of tetrahydrofuran (anhydrous). The solution was  
stirred an additional two hours at 0 °C, filtered, and concentrated to dryness. The  
crude material was purified by silica gel chromatography (230-400 mesh silica gel,  
10 250 g). The column was initially eluted with 20% methanol in methylene chloride,  
then methylene chloride. The crude product was then loaded onto the column and  
eluted with methanol/ methylene chloride/ ammonium hydroxide ( 5/95/0.5 to  
15 15/85/0.5). Product fractions were concentrated to yield 5.28 g (30%) of a thick,  
viscous yellow oil. TLC: 10% methanol in methylene chloride (0.5% ammonium  
hydroxide) on silica gel, R<sub>f</sub> = 0.7. Mass Spectrum (CI) 237.1 (MH<sup>+</sup>, 100%), <sup>1</sup>H  
NMR (200 MHz, CDCl<sub>3</sub>) δ 7.35-7.15 (m, 4H), 4.9 (br. s, 1H), 4.3 (m, 2H), 3.85  
(m, 2H), 1.6 (s, 3H).

##### Amidine Synthesis: Preparation of Intermediate *tert*-butyl 20 N-(3-((acetimidoyl)aminomethyl)benzyl)-carbamate

To a solution of 2.6 g (11.0 mmol) *tert*-butyl N-(3-(aminomethyl) benzyl  
carbamate in 25 ml of absolute ethanol at 0 °C was added 2.22 g (11.0 mmol) of  
intermediate A. The solution was stirred for three hours and concentrated to  
dryness. The residue was partitioned between ether (100 ml) and water (100 ml).  
25 The aqueous was extracted with ether (50 ml) and freeze-dried to yield 3.28 g  
(95%) of white solid carbamate hydrochloride as product. <sup>1</sup>H NMR (200 MHz,  
DMSO) δ 7.4-7.3 (m, 1H), 7.24-7.16 (m, 3H), 4.46 (s, 2H), 4.13 (s, 2H), 2.21  
(s, 3H), 1.4 (s, 9H). Mass Spectrum (CI) 278.1 (MH<sup>+</sup>, 100%).

Deprotection of Intermediate *tert*-butyl N-(3-((Acetimidoyl)aminomethyl)benzyl)-carbamate

5 Solid *tert*-butyl N-(3-((acetimidoyl)aminomethyl)benzyl)carbamate (3.0 g, 9.6 mmol) was suspended into 150 ml anhydrous dioxane. 4N Hydrochloric acid in dioxane solution (20 ml) was added at 20°C. The mixture was stirred for 24 hours at 20°C. After diluting with 200 ml ether, the white solids were collected by filtration and dried under vacuum at 100°C to yield 2.34 g (98%) of N-(3-(aminomethyl)benzyl)-acetamidine. <sup>1</sup>H NMR (200 MHz, D<sub>2</sub>O) δ 7.52-7.35 (m, 4H), 4.49 (s, 2H), 4.16 (s, 2H), 2.25 (s, 3H). Mass Spectrum (CI) 177.9 (MH<sup>+</sup>, 100%).

Example 2

Preparation of N-(3-(Aminomethyl)benzyl)-2-fluoroacetamidine

15 To a solution of 14 g (59.24 mmol) *tert*-butyl N-(3-(aminomethyl)benzyl)carbamate in 100 ml ethanol at 0°C was added 18.8 g (71.1 mmol) of intermediate B. The solution was concentrated after 14 h to a thick oil, stirred vigorously, and 200 ml of ether was added. The residue that separates was washed several times with ether. The resulting solid was stirred in 250 ml of acetic acid and 25 ml of 30% HBr in acetic acid solution. After stirring 30 minutes, 500 ml of ether was added and the solids were collected by filtration and dried to yield 21.55 g (100 %) of N-(3-(Aminomethyl)benzyl)-2-fluoroacetamidine as a white solid. A 2.0 g sample was recrystallized from MeOH/EtOAc to yield 1.43 g of a white solid. Mass spectrum (CI) 196.0 (MH<sup>+</sup>, 76 %). <sup>1</sup>H NMR (200 MHz, DMSO) δ 7.52-7.38 (m, 4H), 5.42 (d, J = 45.3 Hz, 2H), 4.6 (s, 2H), 4.07 (s, 2H).

Example 3

Preparation of N-(4-(Aminomethyl)benzyl)acetamidine

Prepared from *tert*-butyl N-(4-(aminomethyl)benzyl)carbamate and intermediate A

by the method of example 1 except that the crude deprotection product was freeze-dried from 25 ml water. Mass spectrum (CI) 178 ( $MH^+$ , 100 %).  $^1H$  NMR (200 MHz,  $D_2O$ )  $\delta$  7.5-7.35 (m, 4H), 4.5 (s, 2H), 4.15 (s, 2H), 2.15 (s, 3H).

5     **Example 4**

**Preparation of N-(3-(Aminomethyl)benzyl)-2-methoxyacetamidine**

The intermediate *tert*-butyl N-(3-((2-methoxyacetimidoyl)aminomethyl)benzyl)-carbamate (360 mg) prepared from *tert*-butyl N-(3-(aminomethyl)benzyl carbamate and intermediate G analogous to example 1  
10     was deprotected in trifluoroacetic acid at 20 °C for 16 h. The crude product was purified by C18 reverse phase chromatography eluting with water (0.1 % trifluoroacetic acid) to yield 150 mg of N-(3-(Aminomethyl)benzyl)-2-methoxyacetamidine as a yellowed, viscous oil (freeze-dried). Mass spectrum (CI) 208.0 ( $MH^+$ , 100 %).  $^1H$  NMR (200 MHz,  $D_2O$ )  $\delta$  7.5-7.3 (m, 4H), 4.57 (s, 2H), 4.37 (s, 2H), 4.15 (s, 2H), 3.44 (s, 3H).  
15

**Example 5**

**Preparation of N-(3-(Aminomethyl)benzyl)-2-aminoacetamidine**

Prepared from 430 mg of *tert*-butyl N-(3-(aminomethyl)benzyl carbamate and  
20     intermediate I analogous to example 4. The freeze-dried fractions left 640 mg of a yellowed viscous oil. Mass spectrum (CI) 193.1 ( $MH^+$ , 100 %). Capillary electrophoresis produced a single peak.

**Example 6**

25     **Preparation of N-(4-Aminobutyl)acetamidine**

Prepared from 0.98 g (5.23 mmol) N-Boc-1,4-diaminobutane and intermediate B analogous to example 1. The product residue was purified by preparative reverse phase (C18) chromatography eluting with a methanol/water/trifluoroacetic acid gradient (5/95/0.1 to 90/10/0.1) to yield 0.76 g of pale yellow

N-(4-aminobutyl)acetamidine. Capillary electrophoresis gave a single peak (50 mm x 40 cm fused silica capillary, 5.5 minutes). Mass spectrum (CI) 130.0 ( $MH^+$ , 95 %).  $^1H$  NMR (300 MHz,  $D_2O$ )  $\delta$  3.15 (m, 2H), 2.89 (m, 2H), 2.08 (s, 3H), 1.62-1.55 (m, 4H).

5

#### Example 7

##### Preparation of N-(5-Aminopentyl)acetamidine

Prepared from 0.97 g (4.8 mmol) N-Boc-1,5-diaminopentane analogous to example 6. Capillary electrophoresis gave a single peak (50 mm x 40 cm fused silica capillary, 5.9 minutes). Mass spectrum (CI) 144.0 ( $MH^+$ , 100 %).  $^1H$  NMR (300 MHz,  $D_2O$ )  $\delta$  3.12 (m, 2H), 2.85 (m, 2H), 2.08 (s, 3H), 1.6-1.5 (m, 4H), 1.35-1.25 (m, 2H).

10

#### Example 8

##### Preparation of N-(6-Aminohexyl)acetamidine

15

To a 0 °C stirred solution of 1.0 g (4.62 mmol) of N-Boc-1,6-diaminohexane in 30 ml ethanol was added 0.86 g (4.62 mmol) of intermediate B. After two hours the solution was concentrated, diluted with water (50 ml), and extracted with ether twice. The aqueous phase was freeze-dried to yield 0.9 g of white sticky solid tert-butyl N-(6-((acetimidoyl) amino)hexyl)carbamate intermediate. A suspension of the carbamate above (0.88 g, 3 mmol) in 15 ml dioxane and 7.5 ml trifluoroacetic acid was treated with 10 ml (40 mmol) of 4N hydrochloric acid in dioxane solution with stirring overnight. The mixture was diluted with ether (50 ml) and the solids were isolated by filtration to provide 0.7 g N-(6-aminohexyl)acetamidine as a white solid (66 % yield from N-Boc-1,6-diaminohexane). M.P. = 163-164°C. Mass spectrum (CI) 158.1 ( $MH^+$ , 100 %).

20

25

**Example 9****Preparation of N-(7-Aminoheptyl)acetamide**

Prepared from N-Boc-1,7-diaminohexane (1.0 g, 4.34 mmol) by the method of example 8 to yield 0.76 g of the title compound as a white solid. M.P. = 165-168°C. Mass spectrum (CI) 172.1 (MH<sup>+</sup>, 100 %). <sup>1</sup>H NMR (300 MHz, D2O) δ 3.16 (t, J = 7.0 Hz, 2H), 2.9 (t, J = 7.5 Hz, 2H), 2.11 (s, 3H), 1.59-1.5 (m, 4H), 1.28 (br s, 6H).

**Example 10****Preparation of N,N'-(1,3-Phenylenebis(methylene)) bis(acetamide)**

To a solution of 1.0 g (7.34 mmol) m-xylenediamine in 75 ml ethanol at 0 °C was added 3.7 g (15.0 mmol) of solid intermediate B. The solution was stirred at 0 °C for two hours and concentrated. The residue was taken into 50 ml of water and extracted twice with ether (2 x 50 ml). The aqueous solution was freeze-dried to provide a quantitative yield of N,N'-(1,3-phenylenebis(methylene))bis(acetamide) bis-hydrobromide as a white solid (2.81 g). Mass spectrum (CI) 219.0 (MH<sup>+</sup>, 26 %). <sup>1</sup>H NMR (300 MHz, D2O) δ 7.3 (m, 1H), 7.2 (m, 3H), 4.35 (s, 4H), 2.15 (s, 6H).

Examples 11-13 were prepared from the appropriate diamines by methods analogous to example 10:

**Example 11****N,N'-(1,4-Phenylenebis(methylene))bis(acetamide)**

Mass spectrum (FAB) 219.3 (MH<sup>+</sup>, 100 %). <sup>1</sup>H NMR (300 MHz, D2O) δ 7.45 (s, 4H), 4.46 (s, 4H), 2.2 (s, 6H).



**Example 12****N,N'-(1,3-Phenylene(diethylene))bis(acetamidine)**

Mass spectrum (FAB) 247.2 (MH<sup>+</sup>, 100 %). <sup>1</sup>H NMR (200 MHz, D<sub>2</sub>O) δ 7.35-7.15 (m, 4H), 3.50 (t, J = 6.8 Hz, 4H), 2.9 (t, J = 6.9 Hz, 4H), 2.1 (s, 6H).

**Example 13****N,N'-(1,4-Phenylene(diethylene))bis(acetamidine)**

Mass spectrum (CI) 247 (MH<sup>+</sup>, 25 %). <sup>1</sup>H NMR (300 MHz, D<sub>2</sub>O) δ 7.16 (s, 4H), 3.40 (t, J = 6.8 Hz, 4H), 2.81 (t, J = 6.8 Hz, 4H), 2.04 (s, 6H).

**Example 14****Preparation of N-(3-(hydroxymethyl)phenyl)acetamidine**

Prepared from 1.23 g (10 mmol) 3-aminobenzyl alcohol and 2.22 g (11.0 mmol) of intermediate A by the method of example 1 to provide a white solid (2.0 g, 100% yield). Mass spectrum (CI) 165.0 (MH<sup>+</sup>, 100 %). Capillary zone electrophoresis on a fused silica column (75 microns x 40 cm) at pH = 2 gave a 7.78 minute retention time, 99.3 % purity. <sup>1</sup>H NMR (300MHz, D<sub>2</sub>O) δ 7.4 (t, J = 7.7 Hz, 1H), 7.3 (d, J = 7.4 Hz, 1H), 7.18 (s, 1H), 7.13 (d, J = 7.9 Hz, 1H), 4.52 (s, 2H), 2.26 (s, 3H).

**Example 15****Preparation of N-(3-(Aminomethyl)phenyl)acetamidine**

To a stirred solution of 2.5 g (13.25 mmol) 3-nitrobenzylamine hydrochloride and 8.5 ml (30 mmol) of triethylamine in 75 ml anhydrous methanol and 25 ml anhydrous tetrahydrofuran was added 3.19 g (14.5 mmol) di-*tert*butyl dicarbonate. The solution was filtered after 14 hours, concentrated, and purified by silica gel chromatography to yield 3.3 g (100 %) of protected intermediate (silica gel R<sub>f</sub> = 0.45, 30 % ethyl acetate in hexanes). A solution of 3.1 g (12.29 mmol) of this

intermediate was hydrogenated for 90 minutes under 50 psi hydrogen in 75 ml methanol in the presence of 1 g 10 % palladium on carbon. The mixture was filtered and concentrated to an oil (2.65 g). This oil was taken into 30 ml ethanol (0°C) and 2.48 g (12.3 mmol) intermediate A was added. After 3 hours, the solution was concentrated, partitioned between 100 ml ether and 40 ml water, and the aqueous phase was washed with ether (100 ml) and freeze-dried to a white solid (3.5 g, 95% yield). Mass spectrum (CI) 264.1 (MH<sup>+</sup>, 100 %). To a suspension of 2.0 g (6.67 mmol) of this solid in 20 mL dioxane was added 30 ml of 4N hydrochloric acid in dioxane solution. The suspension was stirred for two hours, diluted with 150 ml ether, and filtered. The solids recovered were dried under vacuum to yield 1.36 g (86 %) of the title compound N-(3-aminomethylphenyl)acetamidine dihydrochloride. Mass spectrum (CI) 164.0 (MH<sup>+</sup>, 100 %). Capillary zone electrophoresis on a fused silica column (75 microns x 40 cm) at pH = 2.1 gave a 5.06 minute retention time, 97.6 % purity. <sup>1</sup>H NMR (300MHz, DMSO)  $\delta$  7.6-7.5 (m, 3H), 7.3 (m, 1H), 4.03 (s, 2H), 2.36 (s, 3H).

#### Example 16

##### Preparation of N-[[6-(Aminomethyl)-2-pyridyl]methyl]acetamidine

Prepared from 1.5 g (10.9 mmol) Bis-2,6-(aminomethyl)pyridine ( synthesized by standard methods from 2,6-pyridinedimethanol via bromination with carbon tetrabromide and triphenylphosphine, bis-azide formation with sodium azide in dimethylformamide, and catalytic hydrogenation to the diamine with palladium on carbon) and intermediate B analogous to example 10. The crude product was purified by preparative reverse phase HPLC (C18) eluting with a gradient of methanol/water /trifluoroacetic acid (10/90/0.1 to 90/10/0.1) and freeze-dried to yield N-[[6-(aminomethyl)-2-pyridyl]methyl]acetamidine (clear, colorless, viscous oil, 130 mg). Mass spectrum (CI) 179.0 (MH<sup>+</sup>, 100 %).

**Example 17****Preparation of N-(2-(5-((acetimidoylamino)methyl)-2-thienyl)ethyl)acetamidine**

Prepared from 2-(2-Aminoethyl)-5-(aminomethyl)thiophene and intermediate B analogous to example 10. Mass Spectrum (FAB) 239.1 (MH<sup>+</sup>). <sup>1</sup>H NMR (200 MHz D<sub>2</sub>O) δ 6.95 (d, J = 3.5 Hz, 1H), 6.81 (d, J = 3.4 Hz, 1H), 4.58 (s, 2H), 3.52 (t, J = 6.6 Hz, 2H), 3.1 (s, 3H), 2.14 (s, 3H).

**Example 18****Preparation of N-(3-(5-(2-(Acetimidoylamino)ethyl)-2-thienyl)propyl)acetamidine**

Prepared from 2-(2-Aminoethyl)-5-(3-aminopropyl)thiophene and intermediate B analogous to example 10. Mass Spectrum (FAB) 267.3 (MH<sup>+</sup>). <sup>1</sup>H NMR (200 MHz D<sub>2</sub>O) δ 6.75-6.7 (m, 2H), 3.50 (t, J = 6.6 Hz, 2H), 3.23 (t, J = 6.7 Hz, 2H), 3.07 (t, J = 6.6 Hz, 2H), 2.85 (t, J = 7.3 Hz, 2H), 2.14 (s, 6H), 2.0-1.9 (m, 2H).

**Example 19****Preparation of N-(3-(Aminomethyl)benzyl)-2,2-difluoroacetamidine**

Prepared from tert-butyl N-(3-(aminomethyl)benzyl)carbamate and intermediate F analogous to example 1. <sup>1</sup>H NMR (200 MHz, D<sub>2</sub>O) δ 4.17 (2H, s), 4.64 (2H, s), 6.70 (1H, t, J = 52.5 Hz), 7.47 (4H, m). Mass Spectrum (CI) 214.1 (MH<sup>+</sup>, 100%).

**Example 20****Preparation of N-(3-(Aminomethyl)benzyl)-2,2,2-trifluoroacetamidine**

To an excess of freshly condensed trifluoroacetonitrile stirring at -78 °C was added a solution of tert-butyl N-[3-(aminomethyl) benzyl]carbamate (746 mg, 3.16 mmol) in CH<sub>3</sub>CN (5 ml) dropwise. The mixture froze into a solid white mass. After warming to room temperature, the mixture was warmed gently in a warm

water bath to remove excess trifluoroacetonitrile. The mixture was concentrated at reduced pressure and the residue was chromatographed on silica gel eluting with 2:3 ethyl acetate in hexane to provide 897 mg (86%) of a viscous oil. To a stirred solution of the above oil (887 mg, 2.68 mmol) in dioxane (20 ml) was added 30% HBr in acetic acid (3 ml, 15.0 mmol). The resulting suspension was stirred for 6h, poured into ethyl acetate (200 ml) and stirred for 30 minutes. The solid was collected, suspended in hot iPrOH, cooled to rt. recollected and dried *in vacuo* to afford 731 mg (69%) of N-[3-(aminomethyl)benzyl]-2,2,2-trifluoroacetamide as a dihydrobromide salt. Mp = 242-244°C. 200 MHz <sup>1</sup>H NMR (D<sub>2</sub>O) δ 4.18 (2H, s), 4.67 (2H, s), 7.48 (4H, m).

#### Example 21

##### Preparation of 3-(2-Fluoroacetimidoylamino)benzamide

A solution of 1.0 g (7.34 mmol) of 3-aminobenzamide in 15 ml of absolute ethanol at room temperature was treated with intermediate D and stirred 24 h. After concentration *in vacuo* to 5-10 ml, the residue was treated with Et<sub>2</sub>O and filtered to give 1.69 g of an off-white solid. Recrystallization from MeOH/Et<sub>2</sub>O provided 1.0 g (50%) of the title compound as a white solid: mp 219-220°C; <sup>1</sup>H NMR (D<sub>2</sub>O, 300 MHz) δ 7.70 (d, 1 H, J = 7.6 Hz), 7.66 (s, 1 H), 7.56 (t, 1 H, J = 7.6 Hz), 7.46 (d, 1 H, J = 7.6 Hz), 5.46 (s, 1 H), 5.31 (s, 1 H).

The following compounds (22-43) were prepared in an analogous manner to example 21:

#### Example 22

##### Preparation of 3-(Acetimidoylamino)benzamide

Prepared from 3-aminobenzamide (1.0 g) and intermediate B (1.63 g). The white solids were filtered and dried to yield 1.55 g of 3-(Acetimidoylamino)benzamide. Mass Spectrum (CI) 178.0 (MH<sup>+</sup>, 100%).

**Example 23****Preparation of N-(3-(2-Hydroxyethyl)phenyl)acetamidine**

Prepared from 3-(2-hydroxyethyl)aniline and intermediate A. Mass Spectrum (CI) 179.0 (MH<sup>+</sup>, 100%).

5

**Example 24****Preparation of N-(3-Fluorophenyl)acetamidine**

Prepared from 3-fluoroaniline using intermediate A. Recrystallization from MeOH/Et<sub>2</sub>O provided 816 mg (48%) of the title compound as a white solid: mp 175-177°C; <sup>1</sup>H NMR (D<sub>2</sub>O, 200 MHz) δ 7.52 (m, 1 H), 7.28-7.05 (m, 3 H), 2.37 (s, 3 H).

10

**Example 25****Preparation of 3-(Acetimidoylamino)benzamidine**

Prepared from 3-amino-benzamidino 2HCl using intermediate A and 1 equivalent of Et<sub>3</sub>N. Recrystallization from MeOH/Et<sub>2</sub>O provided 1.48 g (82%) of a white solid: mp 263-265°C; <sup>1</sup>H NMR (D<sub>2</sub>O, 300 MHz) δ 7.13 (dt, 1 H, J = 7.8, 1.5 Hz), 7.66-7.52 (m, 3 H), 2.29 (s, 3H).

15

20

**Example 26****Preparation of N-(3-Nitrophenyl)acetamidine**

Prepared from 3-nitroaniline using intermediate A. Recrystallization from MeOH/Et<sub>2</sub>O gave 513 mg (35%) of a white solid. The solid was dissolved in CH<sub>2</sub>Cl<sub>2</sub>, basified with saturated NaHCO<sub>3</sub>, and extracted with CH<sub>2</sub>Cl<sub>2</sub>. The combined organic layers were dried (Na<sub>2</sub>SO<sub>4</sub>), filtered, and concentrated *in vacuo*. The yellow solid was treated with iPrOH followed by HCl in EtOH. Et<sub>2</sub>O was added, and resulting white solid was collected to give 310 mg of title compound: mp 226-228°C; <sup>1</sup>H NMR (D<sub>2</sub>O, 300 MHz) δ 8.24-8.10 (m, 2 H), 7.70-7.58 (m, 2 H), 2.31 (s, 3 H).

25

**Example 27****Preparation of N-(3-Acetylphenyl)acetamidine**

Prepared from 3-aminoacetophenone using intermediate A. Recrystallization from MeOH/Et<sub>2</sub>O gave 1.16 g (74%) of a white solid. The solid was dissolved in Et<sub>2</sub>O/saturated NaHCO<sub>3</sub>, and the aqueous layer was extracted with Et<sub>2</sub>O. The combined organics were dried (Na<sub>2</sub>SO<sub>4</sub>), filtered, and concentrated *in vacuo*. The residue was treated with Et<sub>2</sub>O and a small amount of HCl/ethanol. The resulting white solid was collected to give 325 mg of title compound: mp 204-206°C; <sup>1</sup>H NMR (D<sub>2</sub>O, 300 MHz) δ 7.92 (dt, 1 H, J = 7.6, 1.2 Hz), 7.77 (br s, 1 H), 7.55 (t, 1 H, J = 7.6 Hz), 7.46 (dt, 1 H, J = 7.6, 1.8 Hz), 2.52 (s, 3 H), 2.29 (s, 3 H).

**Example 28****Preparation of N-(3-Cyanophenyl)acetamidine**

Prepared from 3-aminobenzonitrile using intermediate A. Recrystallization from MeOH/Et<sub>2</sub>O gave 814 mg (50%) of a solid. The solid was dissolved in Et<sub>2</sub>O/saturated NaHCO<sub>3</sub>, and the aqueous layer was extracted with Et<sub>2</sub>O. The combined organics were dried (Na<sub>2</sub>SO<sub>4</sub>), filtered, and concentrated *in vacuo*. The residue was treated with Et<sub>2</sub>O and a small amount of HCl/ethanol. The resulting white solid was collected to give 576 mg of title compound: mp 228-230°C; <sup>1</sup>H NMR (D<sub>2</sub>O, 300 MHz) δ 7.72 (dt, 1 H, J = 7.8, 1.5 Hz), 7.64 (br s, 1 H), 7.62-7.49 (m, 2 H), 2.28 (s, 3 H).

**Example 29****Preparation of N-(3-Bromophenyl)acetamidine**

Prepared from 3-bromoaniline using intermediate A. The resulting solid was dissolved in Et<sub>2</sub>O/saturated NaHCO<sub>3</sub>, and the aqueous layer was extracted with Et<sub>2</sub>O. The combined organics were dried (Na<sub>2</sub>SO<sub>4</sub>), filtered, and concentrated *in vacuo*. The residue was treated with Et<sub>2</sub>O and a small amount of HCl/ethanol.

The resulting white solid was collected to give 776 mg (54%) of title compound: mp 208-210°C; <sup>1</sup>H NMR (D<sub>2</sub>O, 300 MHz) δ 7.52 (d, 1 H, J = 7.8 Hz), 7.43 (br s, 1 H), 7.31 (t, 1 H, J = 7.8 Hz), 7.18 (d, 1 H, J = 7.8 Hz), 2.26 (s, 3H).

5

### Example 30

#### Preparation of N-(3-(Methylthio)phenyl)acetamidine

Prepared from 3-(methylmercapto)-aniline using intermediate A. Recrystallization from MeOH/Et<sub>2</sub>O gave 1.11 g (71%) of a purple solid. The resulting solid was dissolved in Et<sub>2</sub>O/saturated NaHCO<sub>3</sub>, and the aqueous layer was extracted with Et<sub>2</sub>O. The combined organics were dried (Na<sub>2</sub>SO<sub>4</sub>), filtered, and concentrated *in vacuo*. The residue was treated with Et<sub>2</sub>O and a small amount of HCl/ethanol. The resulting solid was collected to give 778 mg of title compound: mp 134-135°C; <sup>1</sup>H NMR (D<sub>2</sub>O, 300 MHz) δ 7.33 (t, 1 H, J = 7.8 Hz), 7.23 (d, 1 H, J = 7.8 Hz), 7.09 (br s, 1 H), 6.97 (d, 1 H, J = 7.8 Hz), 2.36 (s, 3H), 2.25 (s, 3H).

10

15

### Example 31

#### Preparation of N-(3-(1-Hydroxyethyl)phenyl)acetamidine

Prepared from (+)-3-(1-hydroxyethyl)aniline using intermediate C. Upon stirring overnight, the reaction was filtered, and the filtrate was concentrated *in vacuo*. The residue was triturated with absolute EtOH/pentane, and the title compound was collected by filtration to give 2.1 g (88%) of an off-white solid: mp 132-135°C; <sup>1</sup>H NMR (D<sub>2</sub>O, 300 MHz) δ 7.40 (t, 1 H, J = 7.8 Hz), 7.33 (d, 1 H, J = 7.8 Hz), 7.19 (s, 1 H), 7.12 (d, 1 H, J = 7.8 Hz), 4.80 (q, 1 H, J = 6.6 Hz), 2.26 (s, 3 H), 1.32 (d, 3 H, J = 6.6 Hz).

20

25

### Example 32

#### Preparation of N-(3-Acetimidoylamino)benzyl)acetamidine

A solution of 1.2 g (6.36 mmol) of 3-nitrobenzylamineoHCl in 50 ml of EtOH

containing 150 mg of 10% palladium on carbon was placed on a Parr hydrogenator under  $H_2$  for 1h. The reaction was filtered through a pad of celite washing with EtOH, and the filtrate was concentrated *in vacuo*. The residue was dissolved in  $CH_2Cl_2$  and treated with 1.0 N NaOH. The aqueous layer was extracted with  $CH_2Cl_2$ , and the combined organic layers were dried ( $MgSO_4$ ), filtered, and concentrated *in vacuo* to give 600 mg (77%) of 3-aminobenzyl amine which was used directly in the next reaction. The title compound was prepared as in example 21 from 600 mg of 3-aminobenzylamine using 2 equivalents of intermediate C. Recrystallization from MeOH/ $Et_2O$  provided 1.46 g (81%) of a white solid: mp 276-277°C;  $^1H$  NMR ( $D_2O$ , 300 MHz)  $\delta$  7.45 (t, 1 H,  $J = 7.8$  Hz), 7.31 (d, 1 H,  $J = 7.8$  Hz), 7.25-7.15 (m, 2 H), 4.41 (s, 2 H), 2.28 (s, 3 H), 2.15 (s, 3 H).

### Example 33

#### Preparation of N-(3-Acetimidoylamino)benzenesulfonamide

A solution of 1.5 g (7.42 mmol) of 3-nitrobenzenesulfonamide in 50 ml of EtOH containing 150 mg of 10% palladium on carbon was placed on a Parr hydrogenator under  $H_2$  for 1 h. The reaction was filtered through a pad of celite washing with EtOH, and the filtrate was concentrated *in vacuo* to give 1.28 g (100%) of 3-aminobenzenesulfonamide which was used immediately in the next reaction. The title compound was prepared as in example 21 from 3-aminobenzenesulfonamide using intermediate C. Recrystallization from MeOH/ $Et_2O$  provided 1.46 g (67%) of title compound as an off-white solid: mp 240-242°C;  $^1H$  NMR ( $D_2O$ , 300 MHz)  $\delta$  7.86 (d, 1 H,  $J = 8.1$  Hz), 7.75 (s, 1 H), 7.63 (t, 1 H,  $J = 8.1$  Hz), 7.51 (d, 1 H,  $J = 8.1$  Hz), 2.30 (s, 3 H).



Example 34Preparation of N-(3-((1,2,3,4-Tetrahydroisoquinol-2-yl)methyl)phenyl)acetamidine

5 A solution of 1.0 g (4.63 mmol) of 3-nitrobenzyl bromide in 30 ml of toluene was treated with 1.16 ml (9.26 mmol, 2 equiv) of 1,2,3,4-tetrahydroisoquinoline. After heating at reflux for 1 h followed by cooling to 25°C, the reaction was diluted with EtOAc, washed with 1 N NaOH, dried (MgSO<sub>4</sub>), filtered, and concentrated *in vacuo*. The residue was filtered through a pad of silica gel eluting with hexanes/Et<sub>2</sub>O (2:1), and the filtrate was concentrated *in vacuo* to give 1.21 g (76%) of intermediate. 10 A solution of 1.21 g (3.50 mmol) of the above nitroaromatic in 50 ml of EtOAc was treated with 140 mg of 10% palladium on carbon and placed on a Parr hydrogenator under H<sub>2</sub> until gas consumption ceased. The reaction was filtered through a pad of silica gel washing with EtOAc, and the filtrate was concentrated *in vacuo* to give 1.0 g (89%) of a yellow solid that was used immediately in the next reaction. 15 The title compound was prepared from 1.0 g (3.13 mmol) of the above compound as in example 21 using intermediate C. After removal of EtOH, 10 ml of 1,4-dioxane was added followed by 1.2 ml of 3.7 M HBr in AcOH. After stirring for 5 min, the reaction was treated with Et<sub>2</sub>O and stirred 3 h. An orange solid was collected and recrystallized from 20 EtOH/EtOAc to give 366 mg (27%) of title compound as a brown solid: mp 260-262°C; <sup>1</sup>H NMR (D<sub>2</sub>O, 300 MHz) δ 7.64-6.94 (m, 8 H), 4.41 (br s, 2 H), 4.26 (s, 2 H), 3.69 (br s, 1 H), 3.36 (br s, 1 H), 3.06 (br s, 1 H), 2.29 (s, 3 H).

Example 35Preparation of N-(3-(2-Amino-4-thiazolyl)phenyl)acetamidine

25 To a solution of 5 g (20.5 mmol) of α-bromo-3-nitroacetophenone in 20 ml of acetone at 25°C was treated with 1.6 g (20.5 mmol, 1 equiv) of thiourea. After stirring overnight, 6.1 g (98%) of a white solid was collected by filtration. A solution of 5.1 g (16.9 mmol) of the above compound in 200 ml of EtOH was

5 treated with 500 mg of 10% palladium on carbon and placed on a Parr hydrogenator under H<sub>2</sub> for 2.5 h. The reaction was filtered through a pad of celite washing with EtOH, and the filtrate was concentrated in vacuo. The residue was dissolved in CH<sub>2</sub>Cl<sub>2</sub> and basified with 1 N NaOH. The aqueous layer was  
10 extracted with CH<sub>2</sub>Cl<sub>2</sub>, and the combined organic layers were dried (MgSO<sub>4</sub>), filtered, and concentrated *in vacuo* to give 1.78 g (55%) of intermediate which was used directly in the next reaction. The title compound was prepared as in example 21 from 1.03 g (5.38 mmol) of the above compound using intermediate C. After removal of EtOH, the residue was diluted with 10 ml of 1,4-dioxane and  
15 acidified with 3.7 M HBr in AcOH. After stirring 30 min, Et<sub>2</sub>O was added and a solid was collected by filtration. The solid was recrystallized from MeOH/Et<sub>2</sub>O to give 1.09 g (51%) of title compound as a white solid: mp 252-254°C; <sup>1</sup>H NMR (D<sub>2</sub>O, 300 MHz) δ 7.62-7.40 (m, 3 H), 7.29 (br d, 1 H, J = 7.5 Hz), 6.91 (s, 1 H), 2.29 (s, 3 H).

### Example 36

#### Preparation of N-(3-Aminophenyl)acetamide

A solution of 10 g (72.4 mmol) of 3-nitroaniline in 200 ml of CH<sub>2</sub>Cl<sub>2</sub> at 25 °C was treated sequentially with 10 ml (71.4 mmol) of Et<sub>3</sub>N, 17.4 g (72.4 mmol) of BOC<sub>2</sub>O, and a catalytic amount of DMAP. After stirring for 72 h, the reaction  
20 was filtered through a pad of silica gel washing with hexanes/EtOAc (1:1), and the filtrate was concentrated *in vacuo*. Purification by medium pressure silica gel column chromatography using hexanes/EtOAc (3:1 / 1:1) provided 4.4 g (26%) of 3-(N-tert-butoxycarbonyl)-nitroaniline which was used directly in the next  
25 reaction. A solution of 1.5 g (6.3 mmol) of the nitroaromatic in 50 ml of EtOAc was treated with 150 mg of 10% palladium on carbon and placed on a Parr hydrogenator under H<sub>2</sub> for 2 h. The reaction was filtered through a pad of silica gel washing with EtOAc, and the filtrate was concentrated *in vacuo* to give 1.3 g (100%) of intermediate which was used directly in the next reaction. The title

compound was prepared as in example 21 from 1.3 g (6.3 mmol) of 3-(N-tert-butoxycarbonyl)-aminoaniline using intermediate C. After removal of EtOH, the residue was diluted with 10 ml of 1,4-dioxane and treated with 2 ml (7.6 mmol, 1.2 equiv) of 3.7 M HBr in AcOH. After stirring 15 min, Et<sub>2</sub>O was added and a solid was collected by filtration. The solid was recrystallized from MeOH/EtOAc to give 1.15 g (59%) of title compound as a white solid: mp 288-289°C; <sup>1</sup>H NMR (D<sub>2</sub>O, 300 MHz) δ 7.51 (t, 1 H, J = 8.1 Hz), 7.33-7.24 (m, 2 H), 7.20 (t, 1 H, J = 1.8 Hz), 2.27 (s, 3H).

### Example 37

#### Preparation of N-(3-((Methylamino)methyl)phenyl)acetamidine

To a solution of 2.0 g (7.93 mmol) of 3-(N-tert-butoxycarbonyl)-nitrobenzylamine in 32 ml of anhydrous THF at -78°C was added 6.34 ml (9.51 mmol, 1.2 equiv) of 1.5 M LDA in cyclohexane. After stirring for 25 min, 0.592 ml (9.51 mmol, 1.2 equiv) of MeI was added, and the reaction was immediately warmed to 25°C and stirred 4 h. The solution was poured into H<sub>2</sub>O and extracted with Et<sub>2</sub>O. The combined organic layers were dried (MgSO<sub>4</sub>), filtered, and concentrated *in vacuo*. Purification by flash silica gel column chromatography using hexanes/Et<sub>2</sub>O (2:1) provided 2.1 g (59%) of intermediate 3-(N-methyl-N-tert-butoxycarbonyl)nitrobenzylamine. A solution of 1.25 g (4.7 mmol) of the above nitroaromatic in 50 ml of EtOAc at 25°C was treated with 140 mg of 10% palladium on carbon and placed on a Parr hydrogenator under H<sub>2</sub> for 4 h. The reaction was filtered through a pad of silica gel washing with EtOAc. The filtrate was concentrated *in vacuo* to give 1.11 g (100%) of intermediate. The title compound was prepared analogous to example 21 from the above aniline intermediate (1.11 g, 4.69 mmol) and intermediate C. After removal of EtOH, 10 ml of 1,4-dioxane was added followed by 1 ml (5.9 mmol) of 5.9 M HBr/AcOH. After stirring 5 min, Et<sub>2</sub>O was added and stirring was continued for 2 h. Filtration provided a solid that was recrystallized from MeOH/EtOAc to give 950 mg (59%) of title compound as a

white solid: mp 222-223°C; <sup>1</sup>H NMR (D<sub>2</sub>O, 300 MHz) δ 7.54-7.24 (m, 4 H), 4.12 (s, 2 H), 2.60 (s, 3 H), 2.28 (s, 3 H).

### Example 38

#### 5 Preparation of N,N'-(1,3-Phenylene)diacetamidine

Prepared from 1,3-phenylene-diamine using 2 equivalents of intermediate C. Recrystallization from MeOH/Et<sub>2</sub>O provided 1.55 g (53%) of title compound as a white solid: mp 256-258°C; <sup>1</sup>H NMR (D<sub>2</sub>O, 300 MHz) δ 7.56 (t, 1 H, J = 8.1 Hz), 7.36-7.20 (m, 3 H), 2.28 (s, 6 H).

10

### Example 39

#### Preparation of N-(3-Hydroxyphenyl)acetamidine

Prepared from 3-aminophenol using intermediate C. Trituration with MeOH/Et<sub>2</sub>O provided 1.9 g (90%) of title compound as a white solid: mp 115-118°C; <sup>1</sup>H NMR (D<sub>2</sub>O, 300 MHz) δ 7.26 (t, 1 H, J = 8.1 Hz), 6.81 (d, 1 H, 8.1 Hz), 6.73 (d, 1 H, J = 8.1 Hz), 6.66 (br s, 1 H), 2.23 (s, 3 H).

15

### Example 40

#### Preparation of N-(3-(Hydrazinomethyl)phenyl)acetamidine

20 A solution of 1.0 g (5.3 mmol) of 3-nitrophenylhydrazineoHCl in 66 ml of THF was treated with 33 ml of saturated NaHCO<sub>3</sub> followed by 1.15 g (5.3 mmol, 1 equiv) of BOC<sub>2</sub>O. After stirring overnight, the reaction was diluted with H<sub>2</sub>O and extracted with Et<sub>2</sub>O. The combined organic layers were washed with brine, dried (MgSO<sub>4</sub>), filtered, and concentrated *in vacuo* to give 1.33 g (100%) of a yellow solid that was used immediately in the next reaction. A solution of 1.33 g (5.3 mmol) of the crude intermediate in 50 ml of EtOAc at 25°C was treated with 200 mg of 10% palladium on carbon and placed on a Parr hydrogenator under H<sub>2</sub> until gas consumption ceased. The reaction was filtered through a pad of celite washing with EtOAc, and the filtrate was concentrated *in vacuo* to give 1.18 g (100%) of

25

a yellow solid that was used immediately in the next reaction. The title compound was prepared as in example 21 using the above intermediate aniline and intermediate C. After removal of EtOH, 10 ml of 1,4-dioxane was added followed by 1 ml of 5.9 M HBr in AcOH. After stirring for 30 min Et<sub>2</sub>O was added, and  
5 a solid was collected by filtration. The solid was recrystallized from MeOH/EtOAc to give 0.78 g (45%) of title compound as a brown solid: mp 218-220°C; <sup>1</sup>H NMR (D<sub>2</sub>O, 300 MHz) δ 7.38 (t, 1 H, J = 8.1 Hz), 6.96-6.88 (m, 2 H), 6.81 (t, 1 H, J = 2.1 Hz), 2.26 (s, 3 H).

10 **Example 41**

**Preparation of N-(4-(Aminomethyl)phenyl)acetamidine**

A solution of 1.15 g (6.1 mmol) of 4-nitrobenzylamineoHCl in 76 ml of THF was treated with 38 mL of saturated NaHCO<sub>3</sub> followed by 1.33 g (6.1 mmol, 1 equiv) of BOC<sub>2</sub>O. After stirring overnight the reaction was diluted with H<sub>2</sub>O and  
15 extracted with Et<sub>2</sub>O. The combined organic layers were dried (MgSO<sub>4</sub>), filtered, and concentrated *in vacuo* to give 1.54 g (100%) of 4-(N-tert-butoxycarbonyl)-nitrobenzylamine that was used immediately in the next reaction.

A solution of 1.54 g (6.1 mmol) of the above compound in 50 ml of EtOAc was treated with 150 mg of 10% palladium on carbon and placed on a Parr hydrogenator under H<sub>2</sub> under gas consumption ceased. The reaction was filtered through a pad of celite washing with EtOAc, and the filtrate was concentrated *in vacuo* to give 1.36 g (100%) of intermediate aniline. The title compound was  
20 prepared from 1.36 g (6.1 mmol) of the above aniline as in example 21 using intermediate C. After EtOH removal, 10 ml of 1,4-dioxane was added followed by 1.2 ml of 5.9 M HBr in AcOH. After stirring for 5 min Et<sub>2</sub>O was added, and a solid was collected by filtration. The solid was recrystallized from MeOH/EtOAc to give 1.2 g (61%) of title compound as a white solid: mp  
25 256-258°C; <sup>1</sup>H NMR (D<sub>2</sub>O, 300 MHz) δ 7.47 and 7.28 (AB quartet, 2 H, J = 8.4

Hz), 4.10 (s, 2 H), 2.28 (s, 3 H).

#### Example 42

##### Preparation of N-(1,2,3,4-Tetrahydroisoquinolin-7-yl)acetamidine

- 5 A solution of 10.5 g (78.8 mmol) of 1,2,3,4-tetra-hydroisoquinoline in 55 ml of concentrated  $\text{H}_2\text{SO}_4$  at  $0^\circ\text{C}$  was treated with 10 g (98.9 mmol, 1.25 equiv) and warmed to  $25^\circ\text{C}$ . After stirring for 48 h, ice was added followed by  $\text{NH}_4\text{OH}$ . The aqueous layer was extracted with  $\text{CH}_2\text{Cl}_2$ , and the combined organics were dried ( $\text{MgSO}_4$ ), filtered, and concentrated to a volume of 100 ml. The solution
- 10 was treated with 80 ml of 1 M HCl in  $\text{Et}_2\text{O}$  followed by  $\text{Et}_2\text{O}$ . A solid was collected and recrystallized from  $\text{H}_2\text{O}$ /acetone to give 6.0 g (36%) of 7-nitro-1,2,3,4-tetrahydroisoquinolineoHCl. A solution of 1.0 g (4.7 mmol) of 7-nitro-1,2,3,4-tetrahydroisoquinolineoHCl in 58 ml of THF was treated with 29 ml of saturated  $\text{NaHCO}_3$  followed by 1.02 g (4.7 mmol, 1 equiv) of  $\text{BOC}_2\text{O}$ .
- 15 After stirring overnight, the reaction was diluted with  $\text{H}_2\text{O}$  and extracted with  $\text{Et}_2\text{O}$ . The combined organic layers were dried ( $\text{MgSO}_4$ ), filtered, and concentrated *in vacuo*. Purification by flash silica gel column chromatography eluting with hexanes/ $\text{Et}_2\text{O}$  (2:1) provided 1.08 g (83%) of N-tert-butoxycarbonyl-7-nitro-1,2,3,4-tetra hydroisoquinoline. A solution of 1.08
- 20 g (3.88 mmol) of the above nitroaromatic in 50 ml of EtOAc was treated with 100 mg of 10% palladium on carbon and placed on a Parr hydrogenator under  $\text{H}_2$  until gas consumption ceased. The reaction was filtered through a pad of silica gel washing with EtOAc, and the filtrate was concentrated *in vacuo* to give 964 mg of an intermediate aniline that was used immediately in the next reaction. The title
- 25 compound was prepared as in example 21 using 0.964 g (3.88 mmol) of the above aniline and intermediate C. After EtOH removal, 10 ml of 1,4-dioxane was added followed by 0.75 ml of 5.9 M HBr in AcOH. After stirring 5 min  $\text{Et}_2\text{O}$  was added, and a solid was collected. Recrystallization from MeOH/EtOAc provided 530 mg (39%) of title compound as a white solid: mp undefined;  $^1\text{H}$  NMR ( $\text{D}_2\text{O}$ ,

300 MHz)  $\delta$  7.29 (d, 1 H,  $J = 8.4$  Hz), 7.12 (d, 1 H,  $J = 8.4$  Hz), 7.06 (br s, 1 H), 4.27 (s, 2 H), 3.41 (t, 2 H,  $J = 6.6$  Hz), 3.02 (t, 2 H,  $J = 6.6$  Hz), 2.26 (s, 3 H).

5     **Example 43**

**Preparation of N-(1H-Indazol-6-yl)acetamide**

      The title compound was prepared from 6-aminoindazole as in example 21 using intermediate C. Recrystallization from MeOH/EtOAc provided 900 mg (70%) of title compound as a brown solid: mp 244-246°C;  $^1\text{H}$  NMR (DMSO, 300 MHz)  
10      $\delta$  13.30 (s, 1 H), 11.23 (s, 1 H), 9.41 (s, 1 H), 8.53 (s, 1 H), 8.14 (s, 1 H), 7.90 (d, 1 H,  $J = 8.4$  Hz), 7.50 (s, 1 H), 6.98 (d, 1 H,  $J = 8.4$  Hz), 2.34 (s, 3 H).

**Example 44**

**Preparation of N,N'-(3,3'-(Iminodimethylene) diphenyl)diacetamide**

15     A solution of 1.0 g (4.63 mmol) of 3-nitrobenzyl bromide in 20 ml of DMPU was treated with 873 mg (4.63 mmol, 1 equiv) of 3-nitrobenzylamineoHCl followed by 1.5 g (13.9 mmol, 3 equiv) of  $\text{Na}_2\text{CO}_3$ . After stirring at 100°C for 1 h, the reaction was cooled to 25°C and treated with 1.06 g (4.63 mmol, 1 equiv) of  $\text{BOC}_2\text{O}$ . After stirring for 48 h, the reaction was poured into  $\text{H}_2\text{O}$  and extracted  
20     with hexanes/EtOAc (1:1). The combined organic layers were dried ( $\text{MgSO}_4$ ), filtered, and concentrated *in vacuo*. Purification by flash silica gel column chromatography eluting with hexanes/ $\text{Et}_2\text{O}$  (1:1) provided 1.26 g (70%) of bis-nitroaromatic intermediate. A solution of 1.26 g (3.25 mmol) of the above compound in 50 ml of EtOAc was treated with 150 mg of 10% palladium on carbon and placed on a Parr hydrogenator under  $\text{H}_2$  until gas consumption ceased.  
25     The reaction was filtered through a pad of celite washing with EtOAc, and the filtrate was concentrated *in vacuo* to give 1.07 g (100%) of an oil that was used directly in the next reaction. The title compound was prepared from 1.06 g (3.25 mmol) of the above diamine as in Example 21 using 2 equivalents of intermediate

C. After EtOH removal, 10 ml of 1,4-dioxane was added followed by 0.75 ml of 5.9 M HBr in AcOH. After stirring for 5 min Et<sub>2</sub>O was added and stirring was continued for 2 h. A solid was collected and recrystallized from MeOH to give 708 mg (40%) of title compound as a white solid: mp 282-283°C; <sup>1</sup>H NMR (D<sub>2</sub>O, 300 MHz) δ 7.54-7.25 (m, 8 H), 4.23 (s, 4 H), 2.28 (s, 6 H). The following compound (45) was prepared in an analogous fashion:

#### Example 45

##### Preparation of N-(3-((Hydroxyamino)methyl)phenyl)acetamidine

The title compound was prepared from 482 mg (6.94 mmol) of hydroxylamine and intermediate C for the amidine formation. Recrystallization from MeOH/EtOAc provided 600 mg (47%) of title compound as a white solid: mp 214-215°C; <sup>1</sup>H NMR (D<sub>2</sub>O, 300 MHz) δ 7.54-7.25 (m, 2 H), 7.36-7.28 (m, 2 H), 4.35 (s, 2 H), 2.28 (s, 3 H).

#### Example 46

##### Preparation of N-(3-(1H-Pyrazol-3-yl)phenyl)acetamidine

A solution of 5.0 g (30.3 mmol) of 3-nitroacetophenone in 16 ml of DMF was treated with 4.8 mL of (MeO)<sub>2</sub>CHNMe<sub>2</sub> and heated to 120°C for 24 h. After cooling to 25°C, the reaction poured into 1 N HCl and extracted with hexanes/EtOAc (1:1). The combined organic layers were washed with brine, dried (MgSO<sub>4</sub>), filtered, and concentrated in vacuo to give 1.88 g (28%) of intermediate.

A solution of 940 mg (4.27 mmol) of the above compound in 34 ml of MeOH and 8.5 ml of H<sub>2</sub>O was treated with 0.855 ml (17.6 mmol) of H<sub>2</sub>NNH<sub>2</sub>·O and heated at reflux for 2 h. After cooling to 25°C, the reaction was poured into 1 N HCl and extracted with EtOAc. The combined organic layers were dried (MgSO<sub>4</sub>), filtered, and concentrated *in vacuo* to give 808 mg (100%) of 1-(1H-pyrazol-3-yl)-3-nitrobenzene.

A solution of 808 mg (4.27 mmol) of the above intermediate in 21 ml of



$\text{CH}_2\text{Cl}_2$  was treated with a catalytic amount of DMAP followed by 918 mg (4.27 mmol, 1 equiv) of  $\text{BOC}_2\text{O}$ . After stirring overnight, the reaction was concentrated *in vacuo* and used immediately in the next reaction.

5 A solution of 1.23 g (4.27 mmol) of the above intermediate in 50 ml of EtOAc was treated with 150 mg of 10% palladium on carbon and placed on a Parr hydrogenator under  $\text{H}_2$  atmosphere until gas consumption ceased. The reaction was filtered through a pad of celite washing with EtOAc, and the filtrate was concentrated *in vacuo* to give an oil that was used immediately in the next reaction.

10 The title compound was prepared from 1.1 g (4.27 mmol) of the above intermediate as in Example 21 using intermediate C. After EtOH removal, 10 ml of 1,4-dioxane was added followed by 0.80 ml of 5.9 M HBr in AcOH. After stirring for 30min  $\text{Et}_2\text{O}$  was added, and stirring was continued for 2 h. A solid was collected and recrystallized from EtOH/EtOAc/ $\text{Et}_2\text{O}$  to give 381 mg (25%) of  
15 title compound as a brown solid: mp 224-226°C;  $^1\text{H}$  NMR ( $\text{D}_2\text{O}$ , 300 MHz)  $\delta$  7.81 (d, 1 H,  $J = 2.4$  Hz), 7.67 (dd, 1 H,  $J = 8.1, 1.2$  Hz), 7.55 (br s, 1 H), 7.49 (t, 1 H,  $J = 8.1$  Hz), 7.25 (br d, 1 H,  $J = 8.1$  Hz), 6.75 (d, 1 H,  $J = 2.4$  Hz), 2.28 (s, 3 H).

20 Example 47

Preparation of N-(3-(Morpholinomethyl)phenyl)acetamidine 2HBr:

A solution of 1.0 g (4.63 mmol) of 3-nitrobenzyl bromide in 30 ml of toluene was treated with 0.806 ml (9.26 mmol, 2 equiv) of morpholine and heated to reflux for 3 h. After cooling to 25°C EtOAc was added, and the organic layer was washed  
25 with 1 N NaOH, dried ( $\text{MgSO}_4$ ), filtered, and concentrated *in vacuo* to give 1.03 g (100%) of an oil.

The crude oil from above was dissolved in 50 ml of EtOAc, treated with 100 mg of palladium on carbon, and placed on a Parr hydrogenator under  $\text{H}_2$  for 1.5 h. The mixture was filtered through a pad of silica gel washing with

EtOAc/MeOH (10:1), and the filtrate was concentrated in vacuo to give 900 mg of an oil.

The title compound was prepared as in Example 21 from 890 mg (4.63 mmol) of the above oil using intermediate C. After EtOH removal, 10 ml of 1,4-dioxane was added followed by 0.90 ml of 5.9 M HBr in AcOH. After stirring for 5 min Et<sub>2</sub>O was added, and a solid was collected. Recrystallization from EtOH/EtOAc provided 938 mg (51%) of title compound as a white solid: mp 219-220°C; <sup>1</sup>H NMR (D<sub>2</sub>O, 300 MHz) δ 7.56-7.28 (m, 4 H), 4.28 (s, 2 H), 3.93 (br s, 2H), 3.63 (br s, 2 H), 3.40-3.00 (m, 4 H), 2.28 (s, 3 H).

The following compounds (48-50) were made in an analogous fashion:

#### Example 48

##### Preparation of N-(3-(1H-Pyrazol-1-ylmethyl)phenyl)acetamidino2 HBr

The title compound was prepared from pyrazole as in Example 47 utilizing intermediate C for the amidine formation. Recrystallization from MeOH/EtOAc provided 960 mg (55%) of a white solid: 218-220°C; <sup>1</sup>H NMR (D<sub>2</sub>O, 300 MHz) δ 7.79 (s, 1 H), 7.63 (s, 1 H), 7.40 (t, 1 H, J = 7.8 Hz), 7.24-7.13 (m, 2 H), 7.03 (s, 1 H), 6.38 (m, 1 H), 5.35 (s, 2 H), 2.23 (s, 3 H).

#### Example 49

##### Preparation of N-(3-((Dimethylamino)methyl)phenyl) acetamidino2HBr

The title compound was prepared from 2 M HNMe<sub>2</sub> in THF as in Example 47 utilizing intermediate C for the amidine formation. Recrystallization from MeOH/EtOAc provided 960 mg (56%) of a white solid: mp 242-243 °C; <sup>1</sup>H NMR (D<sub>2</sub>O, 300 MHz) δ 7.58-7.28 (m, 4 H), 4.23 (s, 2 H), 2.73 (s, 6 H), 2.28 (s, 3 H); low resolution MS (CI) 192 (MH<sup>+</sup>); Anal. Calcd. for C<sub>11</sub>H<sub>17</sub>N<sub>3</sub>O<sub>2</sub>HBr: C, 37.42; H, 5.42; N, 11.90; Br, 45.26. Found: C, 37.45; H, 5.46; N, 11.81; Br, 45.16.

**Example 50****Preparation of N-(3-((2-pyridylamino)methyl) phenyl)acetamidino2HBr**

5 A solution of 1.0 g (4.63 mmol) of 3-nitrobenzyl bromide in 30 ml of toluene was treated with 872 mg (9.26 mmol, 2 equiv) of 2-aminopyridine and heated to reflux for 4 h. Upon cooling to 25°C the reaction mixture was basified with 1 N NaOH and extracted with EtOAc. The combined organic layers were dried (MgSO<sub>4</sub>), filtered, and concentrated *in vacuo* to an oil that was used in the next reaction.

10 A solution of 1.06 g (4.63 mmol) of the above amine in 30 ml of THF was treated with 1.5 g (6.95 mmol, 1.5 equiv) of BOC<sub>2</sub>O and stirred 16 h. The reaction was treated with 1.5 g (6.95 mmol, 1.5 equiv) of BOC<sub>2</sub>O and stirred an additional 7.5 h. The reaction was diluted with saturated NaHCO<sub>3</sub> and extracted with Et<sub>2</sub>O. The combined organic layers were dried (MgSO<sub>4</sub>), filtered, and concentrated *in vacuo*. Purification by flash silica gel column chromatography eluting with hexanes/EtOAc (3:1) provided 411 mg (27%) of an oil that was used

15 in the next reaction.

A solution of 411 mg (1.25 mmol) of the above nitroaromatic in 20 ml of EtOAc was treated with 100 mg of 10% palladium on carbon and placed on a Parr hydrogenator under H<sub>2</sub> for 3 h. The reaction was filtered through celite washing with EtOAc, and the filtrate was concentrated *in vacuo* to give 374 mg (100%) of an oil.

20

The title compound was prepared from 374 mg (1.25 mmol) of the above oil as in example 47 using intermediate C. After removal of EtOH, 5 ml of 1,4-dioxane was added following by 0.30 ml of 5.9 M HBr in AcOH. After stirring for 5 min Et<sub>2</sub>O was added, and stirring was continued overnight. A solid

25 was collected and recrystallized from MeOH/EtOAc to give 270 mg (54%) of title compound as a white solid: mp 247-248 °C; <sup>1</sup>H NMR (D<sub>2</sub>O, 300 MHz) δ 7.76 (ddd, 1 H, J = 9.2, 7.2, 1.8 Hz), 7.65 (d, 1 H, J = 6.3 Hz), 7.41 (t, 1 H, J = 7.6 Hz), 7.33 (d, 1 H, J = 7.6 Hz), 7.22-7.13 (m, 2 H), 6.88 (d, 1 H, J = 9.2 Hz), 6.77 (t, 1 H, J = 6.3 Hz), 4.53 (s, 2 H), 2.25 (s, 3 H).

**Example 51****Preparation of S-(3-(Acetimidoylamino)benzyl)isothioureao2HBr**

A solution of 1.0 g (4.63 mmol) of 3-nitrobenzyl bromide in 10 ml of acetone was treated with 352 mg (4.63 mmol, 1 equiv) of thiourea and stirred for 3 h. The reaction was filtered to give 1.29 g (96%) of a white solid that was used directly in the next reaction.

A solution of 1.29 g (4.41 mmol) of the above compound in 50 ml of EtOH was treated with 200 mg of 10% palladium on carbon and placed on a Parr hydrogenator under H<sub>2</sub> atmosphere overnight. The reaction mixture was filtered through a pad of celite washing with EtOH, and the filtrate was concentrated *in vacuo* to give 1.26 g (100%) of an orange solid that was used immediately in the next reaction.

The title compound was prepared from 1.16 g (4.41 mmol) of the above intermediate as in Example 21 using intermediate C. Recrystallization from MeOH/EtOAc provided 672 mg (40%) of title compound as a brown solid: mp 228-230°C; <sup>1</sup>H NMR (D<sub>2</sub>O, 300 MHz) δ 7.46-7.12 (m, 4 H), 4.31 (s, 2 H), 2.27 (s, 3 H).

**Example 52****Preparation of N-(3-(Aminomethyl)phenyl)-2-fluoroacetamidine****Preparation of intermediate Tert-butyl N-(3-nitro) benzylcarbamate**

To a 0°C solution of 30 g (0.159 mol) of 3-nitrobenzylamine hydrochloride in MeOH (100 ml), THF (100 ml) and 48.8 ml triethylamine was added a solution of 38.2 g (0.175 mol) di-*tert*-butyl dicarbonate in THF (200 ml) over 60 min. After 1 hr, the solution was concentrated, diluted with 100 ml water, and extracted with EtOAc (2 x 100 ml). The dried (Na<sub>2</sub>SO<sub>4</sub>) solution was concentrated and purified by silica gel chromatography eluting with 10% - 30% EtOAc in hexanes to yield 38 g (95%) of *tert*-butyl N-(3-nitro)benzylcarbamate.

Amidine synthesis:

A solution of 10 g (39.64 mmol) of *tert*-butyl N-(3-nitro)benzyl -carbamate in 100 ml EtOAc was hydrogenated under 55 psi H<sub>2</sub> in the presence of 1 g of 10 % Pd on carbon for 90 min. The residue after filtration and concentration was taken into 100 ml of absolute ethanol at 0°C and 16 g (50.8 mmol) of intermediate E was added. The solution was stirred for 16 h and concentrated to a volume of 15 ml before 500 ml of ether was added with rapid stirring. The resulting residue was taken into 75 ml of acetic acid and 15 ml of 30 % HBr in acetic acid solution was added. After 15 min 500 ml of ether was added, the mixture stirred for 30 min, and the pale yellow solids collected via filtration to yield 13.5 g (99 %). Recrystallization of 7 g from 30 ml MeOH and 150 mL EtOAc gave 6.0 g of white solid N-(3-(Aminomethyl)phenyl) -2-fluoroacetamidine. Mass Spectrum (CI)182 (MH<sup>+</sup>, 9.7 %). <sup>1</sup>H NMR (300 MHz, DMSO) δ 7.61-7.32 (m, 4H), 5.5 (d, J = 45.1 Hz, 2H), 4.08 (s, 2H).

Example 53Preparation of N-(3-(2-Aminoethyl)phenyl)acetamidine

3-Nitrophenethyl alcohol (2.5g) was hydrogenated (55 psi, 20°C) in MeOH (100 ml) in the presence of di-*tert*-butyl dicarbonate (3.6 g) and 750 mg of 10 % palladium on carbon to give 3.7 g (100 %) of BOC-protected intermediate (mass spec. 238.0, MH<sup>+</sup>). 3.55 g of this alcohol intermediate was converted to azide via the bromide (CBr<sub>4</sub>, PPh<sub>3</sub>, CH<sub>2</sub>Cl<sub>2</sub>, then NaN<sub>3</sub> in DMF) to provide 1.65 g. Reduction of the azide (1.16 g, EtOAc, Pd/C, 20 °C, 55 psi H<sub>2</sub>) followed by protection (K<sub>2</sub>CO<sub>3</sub>, CH<sub>2</sub>Cl<sub>2</sub>, H<sub>2</sub>O, CBz-Cl) gave 1.6 g (100 %) of intermediate. The BOC protecting group was removed at 0 °C in CH<sub>2</sub>Cl<sub>2</sub> with TFA. The crude material was free based with NaHCO<sub>3</sub> and the resulting residue (with intermediate A) was converted to the intermediate N-(3-(2-(Benzyloxycarbonyl)aminoethyl)phenyl)acetamidine (1.5 g) analogous to example 1. A MeOH (75 ml) and 3.0 M HCl in EtOH (25 ml) solution of this acetamidine (1.2 g) was

hydrogenated at 55 psi H<sub>2</sub> at 20°C for 4h. Filtration through celite and concentration gave the title compound. Mass Spectrum (CI) 178.2 (MH<sup>+</sup>, 100 %). <sup>1</sup>H NMR (300 MHz, D<sub>2</sub>O) δ 7.4-7.1 (m, 4H), 3.14 (t, J = 7.1 Hz, 2H), 2.89 (t, J = 7.1 Hz, 2H), 2.25 (s, 3H).

5

#### Example 54

##### Preparation of N-(3-(Aminomethyl)benzyl)propionamidine

Prepared from *tert*-butyl N-(3-(aminomethyl)benzyl) carbamate (1.0 g) and intermediate J as in example 2 to yield 1.25 g of white solid. A 1.0 g portion of this solid was recrystallized from MeOH/EtOAc to yield 0.93 g of the title compound. <sup>1</sup>H NMR (200 MHz, D<sub>2</sub>O) δ 7.5-7.38 (m, 4H), 4.52 (s, 2H), 4.18 (s, 2H), 2.53 (q, J = 7.7 Hz, 2H), 1.25 (t, J = 7.7 Hz, 3H).

10

#### Example 55

##### Preparation of 2-Amino-N-(3-(Aminomethyl)phenyl)acetamidine

Prepared from *tert*-butyl N-(3-(aminomethyl)benzyl)carbamate and intermediate I analogous to example 2. <sup>1</sup>H NMR (200 MHz, D<sub>2</sub>O) δ 4.25 (4H, d, J=5.5Hz), 7.78-7.40 (4H, m). Mass Spectrum (CI) 179.1 (MH<sup>+</sup>, 100%).

15

20

#### Example 56

##### Preparation of N-(3-(Aminomethyl)phenyl)-2-(methylthio)acetamidine

Prepared from *tert*-butyl N-(3-(aminomethyl)benzyl)carbamate and intermediate H analogous to example 1. <sup>1</sup>H NMR (200 MHz, DMSO) δ 2.30 (3H, s), 3.64 (2H, s), 4.12 (2H, s), 7.65-7.32 (4H, m). Mass Spectrum (CI) 210.0 (MH<sup>+</sup>, 60.5%).

25

#### Example 57

##### Preparation of N-(1,2,3,4-tetrahydro-7-isoquinolyl)methyl)acetamidineo2HBr

To a solution of 5.0 g (37.5 mmol) of 1,2,3,4,-tetra-hydroisoquinoline in 23 ml of concentrated H<sub>2</sub>SO<sub>4</sub> at 25 °C was treated with 6.7 g (37.5 mmol, 1 equiv) of

N-hydroxymethylphthalimide and stirred for 7 days. The reaction was poured into H<sub>2</sub>O, basified with solid NaHCO<sub>3</sub>, and extracted with Et<sub>2</sub>O. The combined organics were dried (MgSO<sub>4</sub>), filtered, and concentrated *in vacuo*. The residue was dissolved in Et<sub>2</sub>O and acidified with 1 M HCl in Et<sub>2</sub>O. The solid was collected and recrystallized from MeOH/Et<sub>2</sub>O to give 1.7 g (10%) of intermediate that was used in the next reaction.

A solution of 500 mg (1.52 mmol) of the above compound in 19 ml of THF was treated with 9 ml of saturated NaHCO<sub>3</sub> followed by 332 mg (1.52 mmol, 1 equiv) of BOC<sub>2</sub>O. After stirring for 48 h, the reaction was diluted with H<sub>2</sub>O and extracted with Et<sub>2</sub>O. The combined organic layers were dried (MgSO<sub>4</sub>), filtered, and concentrated *in vacuo*. Purification by flash silica gel column chromatography eluting with hexanes/Et<sub>2</sub>O (1:1) provided 474 mg (79%) of N-protected tetrahydroisoquinoline.

A solution of 474 mg (1.21 mmol) of the above compound in 12 ml of EtOH was treated with 0.117 ml (2.41 mmol, 2 equiv) of H<sub>2</sub>NNH<sub>2</sub>·OH<sub>2</sub>O and stirred for 36 h. The reaction was diluted with CH<sub>2</sub>Cl<sub>2</sub> and filtered. After concentration *in vacuo*, the residue was dissolved in CH<sub>2</sub>Cl<sub>2</sub> and filtered. The filtrate was concentrated *in vacuo* and used in the next reaction.

The title compound was prepared from the above intermediate as in example 21 using intermediate C. After EtOH removal, 2 ml of 1,4-dioxane was added followed by 0.50 ml of 5.9 M HBr in AcOH. After stirring for 15 min Et<sub>2</sub>O was added, and the supernatant was decanted. The residue was triturated with isopropanol overnight. Filtration provided 272 mg (62%) of title compound as an off-white solid: mp 215-218 °C; <sup>1</sup>H NMR (D<sub>2</sub>O, 300 MHz) δ 7.12-6.88 (m, 3 H), 4.21 (s, 2 H), 4.12 (s, 2 H), 3.27 (t, 1 H, J = 6.3 Hz), 2.87 (t, 1 H, J = 6.3 Hz), 2.01 (s, 3 H).

**Example 58****Preparation of 2-Fluoro-N-(3-(2-hydroxymethyl)phenyl)acetamide**

The title compound (2.19 g) was prepared as a pale yellow solid from 3-aminophenethyl alcohol (1.0 g) and intermediate D (2.32 g) as in example 1.

5  $^1\text{H}$  NMR (300 MHz,  $\text{D}_2\text{O}$ )  $\delta$  7.5 (t,  $J = 7.5$  Hz, 1H), 7.37 (d,  $J = 7.8$  Hz, 1H), 7.2 (m, 2H), 5.4 (d,  $J = 44.9$  Hz, 2H), 3.83 (t,  $J = 6.4$  Hz), 2.87 (t,  $J = 6.4$  Hz, 2H). Mass Spectrum (CI) 197.0 (MH $^+$ , 100 %)

**Example 59**

10 **Preparation of N-(3-Amino-5-indanyl)acetamide**

6-Nitroindan-1-one (18.1 g, 102 mmol) was dissolved in ethanol (400 ml) and reduced using Pd (5%) on carbon (2.0 g) and 35 psi hydrogen pressure on a Parr apparatus. Filtration on celite followed by evaporation gave 6-aminoindan-1-one (13.7 g, 93 mmol). NMR (DMSO) 7.2 (d,  $J = 7.7$  Hz, 1H), 6.9 (dd,  $J_{ab} =$   
15 7.7 Hz,  $J_{ax} = 2.3$  Hz, 1H), 6.75 (d,  $J = 2.3$  Hz, 1H), 5.3 (brs, 2H), 2.9 (t,  $J = 5.6$  Hz, 2H), 2.55 (t,  $J = 5.6$  Hz, 2H).

6-Aminoindan-1-one (10.0 g, 68 mmol) was treated with pyridine (50 ml) and hydroxylamine-O-benzyl ether.HCl (11.4 g, 71.3 mmol) and heated to 50 °C  
20 under  $\text{N}_2$  for 40 min. The mixture was evaporated *in vacuo* and diluted with  $\text{CH}_2\text{Cl}_2$  (300 ml). Filtration gave a solid which was washed with a small volume of  $\text{CH}_2\text{Cl}_2$  and dried *in vacuo* to give 6-Amino-1-O-benzyloximinoindane (14.0 g, 48.5 mmol) as the HCl salt. NMR (DMSO) 10.5 - 9.0 (br s, 3H), 7.5 - 7.3 (m, 8H), 5.2 (s, 2H), 3.05 (m, 2H), 2.9 (m, 2H).

25

6-Amino-1-O-benzyloximinoindane.HCl (5.0 g, 17.3 mmol) was treated with 1.0 M BH3-THF (86 ml) and warmed under  $\text{N}_2$  to 40 °C for 23 hrs. The mixture was evaporated and diluted with  $\text{CH}_2\text{Cl}_2$  and quenched by dropwise addition of MeOH. The solvent was evaporated and 1.0 N HCl (50 ml) was added and the



5 mixture warmed to 50°C for 1hr. After cooling, 1.0 N NaOH (50 ml) was added and the pH adjusted to 9.5 with additional 1.0 N NaOH. The mixture was extracted with CH<sub>2</sub>Cl<sub>2</sub> and the extract evaporated onto silica gel and chromatographed using CH<sub>2</sub>Cl<sub>2</sub> with 30 volume % MeOH. Evaporation of appropriate fractions gave 1,6-diaminoindane (2.0 g, 78 %). NMR (DMSO-d<sub>6</sub>) 6.8 (d, J = 6.8 Hz, 1H), 6.57 (s, 1H), 6.4 (d, J = 6.8 Hz, 1H), 4.8 (brs, 2H), 4.0 (t, J = 7Hz, 1H), 2.75-2.6 (m, 2H), 2.3-2.2 (m, 1H), 1.8 (br s, 2H, NH<sub>2</sub>), 1.6-1.4 (m, 1H).

10 1,6-Diaminoindane ( 0.5 g, 3.4 mmol) was stirred in CH<sub>2</sub>Cl<sub>2</sub> (10ml) and cooled to 0°C. Di-tert-butyl dicarbonate (0.74 g, 3.4 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (3 ml) was added dropwise over 10 min and the solution stirred overnight. The solution was evaporated *in vacuo* and diluted with EtOH (20 ml) and cooled to 0 °C. Intermediate C was added and the mixture allowed to warm to 20 °C and stirred for 90 min. Dropwise addition of 30% HBr / HOAc (1.5 ml) with stirring gave after 70 min a white solid which was collected and washed with a small volume of CH<sub>2</sub>Cl<sub>2</sub> and dried *in vacuo* yielding N-(3-Amino-5-indanyl)acetamidine (1.1 g, 93 %) as the 2HBr salt. NMR (DMSO-d<sub>6</sub>) 8.5 (brs, NH), 8.4 (brs, NH), 7.55 (s, 1H), 7.5 (s, 1H), 7.3 (d, J = 7.0 Hz, 1H), 4.8 (m, 1H), 3.8 (brs, NH), 3.2-3.0 (m, 1H), 3.0-2.8 (m, 1H), 2.6-2.5 (m, 1H), 2.4 (s, 3H, CH<sub>3</sub>), 2.1-2.0 (m, 1H).

### Example 60

#### Preparation of N-(1-Amino-4-indanyl)acetamidine

25 4-Nitroindan-1-one was converted to the 1-oximinobenzylether and reduced to 1-amino-4-nitroindane using BH<sub>3</sub> - THF by the methods described in example 59.

1-Amino-4-nitroindane (0.4 g, 2.24 mmol) was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (10 ml) and treated with t-BOC anhydride (0.59 g, 2.7 mmol). After 18 hrs the solvent was

removed *in vacuo* and the residue dissolved in EtOH (50 ml) and reduced using 5% Pd/C (0.2 g) and H<sub>2</sub> at 32 psi on a Parr apparatus. Evaporation of the solvent to a final volume of 20 ml followed Filtration on celite. The intermediate was treated with intermediate C (0.68g, 2.3 mmol) and stirred for 2 hrs. The solvent  
5 was removed *in vacuo* and the residue dissolved in dioxane (15 ml) and treated with 30% HBr - HOAc. After 90 min the crystalline solid that resulted was collected and washed with a small volume of CH<sub>2</sub>Cl<sub>2</sub> and dried *in vacuo* to give N-(1-Amino-4-indanyl)acetamidine (0.60g, 99%) as the HBr salt. NMR (D<sub>2</sub>O) 7.6 (d, J = 7 Hz, 1H), 7.5 (dd, J = 7 Hz, 1H), 7.3 (d, J = 7Hz, 1H), 4.9 (m, 1H),  
10 3.1-2.8 (m, 2H), 2.6 (m, 1H), 2.4 (s, 3H, CH<sub>3</sub>), 2.1 (m, 1H).

### Example 61

#### Preparation of N-(6-(Aminomethyl)-1-indanyl)acetamidine

6-Cyanoindan-1-one (prepared using the procedure of Allinger and Jones, JOC  
15 1962, p70. vol 27 for the tetralone analogue) (3.86 g, 24.5 mmol) was converted to the 1-oxime-O-benzyl ether and reduced with BH<sub>3</sub>-THF by the methods described in example 59 to give 1-amino-6-(aminomethyl)indane. The diamine was isolated in 31% yield as the 2HCl salt by treatment of an ether solution with 4.0 N HCl in dioxane. 1-Amino-6-aminomethylindane 2HCl (0.5 g, 2.1 mmol) was  
20 stirred in EtOH (20 ml), cooled to 0 °C and treated with NEt<sub>3</sub> (0.59 ml, 4.2 mmol). Intermediate A (0.43 g, 2.1 mmol) was added and the mixture was allowed to warm to 20°C and stir overnight. The white crystalline product was collected and washed with a small volume of ether and dried *in vacuo* to give N-(6-(Aminomethyl)-1-indanyl)acetamidine (0.27 g, 35 %) as the 2HCl salt. NMR  
25 (D<sub>2</sub>O) 7.45-7.25 (m, 3H), 4.8 (m, 1H), 4.45 (s, 2H), 3.2-2.85 (m, 2H), 2.6(m, 1H), 2.2 (s, 3H, CH<sub>3</sub>), 2.1 (m, 1H).

**Example 62****Preparation of N-((3-Amino-5-indanyl)methyl)acetamidine**

1-Amino-6-aminomethylindane . 2 HCl (2.0 g, 8.5 mmol) was stirred in MeOH and cooled to 0°C and treated with Et<sub>3</sub>N (1.8 ml, 12.9 mmol). Di-*tert*-Butyl dicarbonate (1.0 g, 4.6 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (3 ml) was added dropwise over 10 min and the mixture allowed to warm to 20°C and stirred for 60 min. The mixture was evaporated in vacuo followed by treatment with CH<sub>2</sub>Cl<sub>2</sub> (15 ml) gave a white solid which was collected and dried *in vacuo* to yield 1-amino-6-*tert*-butylcarbamoylmethylindane HCl (0.46 g, 18 %). NMR (DMSO) 8.3 (br s, NH<sub>3</sub>), 7.4 (br s, 2H, NHCO), 7.25 (d, J = 7.6 Hz, 1H), 7.2 (d, J = 7.6 Hz, 1H), 4.7 (t, J = 6 Hz, 1H), 4.13 (d, J = 6 Hz, 2H), 3.15-2.95 (m, 1H), 2.95-2.75 (m, 1H), 2.5-2.4 (m, 1H), 2.1-1.9 (m, 1H), 1.4 (s, 9H).

1-Amino-6-*tert*-butylcarbamoylmethylindane hydrochloride (0.15 g, 0.57 mmol) was stirred in EtOH (3 ml) at 0°C and treated with Et<sub>3</sub>N (0.08 ml, 0.57 mmol). Intermediate A (0.115 g, 0.57 mmol) was added and the mixture allowed to warm to 20°C and stir overnight. The mixture was evaporated *in vacuo* and the residue dissolved in dioxane (3 ml) and treated dropwise with 4.0 N HCl/dioxane (1.0 ml). On standing at 5 °C overnight, a white crystalline solid formed and was collected and washed with a small volume of cold dioxane and dried *in vacuo* to give N-((3-Amino-5-indanyl)methyl)acetamidine (0.10 g, 63 %) as the 2HCl salt. NMR (D<sub>2</sub>O) 7.4-7.3 (m, 3H), 5.2 (m, 1H), 4.1 (s, 2H), 3.1-2.8 (m, 2H), 2.6 (m, 1H), 2.2 (s, 3H, CH<sub>3</sub>), 2.0 (m, 1H).1H), 3.0-2.8 (m, 1H), 2.6-2.5 (m, 1H), 2.4 (s, 3H, CH<sub>3</sub>), 2.1-2.0 (m, 1H).

**Example 63****Preparation of 1-Amino-4-(1-iminoethylamino)but-2-yne 2.5hydrochloride****Preparation of 1-Azido-4-(butoxycarbonylamino)but-2-yne**

Hexamethylenetetramine (28.27g) and 1,4-dichlorobut-2-yne (24.84g) were

refluxed in chloroform (165ml) for 20min. Upon cooling to room temperature the resulting precipitate was filtered off and sucked dry. The mother liquor was concentrated to one quarter volume and a further crop of precipitate obtained. Total yield of complex = 46.10g.

5

The above complex (6.7g) was added to a mixture of ethanol (100ml) and concentrated hydrochloric acid (9.4ml) and the mixture refluxed for 3h. On cooling, the precipitate was filtered and discarded. The filtrate was concentrated until it became cloudy at which point it was diluted with diethyl ether (4 volumes).

10

The resulting precipitate was filtered, washed with more diethyl ether and sucked dry. The filtrate was concentrated to a small volume, more diethyl ether added and the resulting precipitate filtered off and sucked dry. The total yield of 1-amino-4-chlorobut-2-yne hydrochloride was 2.924g.

15

1-Amino-4-chlorobut-2-yne hydrochloride (6g) and di-*t*-butyldicarbonate (9.354g) in dry THF (60ml) were treated with diethylisopropylamine (7.52ml) and the mixture stirred overnight at room temperature. After this time the precipitate was filtered off and the filtrate concentrated *in vacuo*. The residue was partitioned between diethyl ether and water and the organic layer washed successively with 0.1N HCl, water and brine, and dried over MgSO<sub>4</sub>. Concentration *in vacuo* gave a green oil (9.205g) which was used directly. The oil was dissolved in DMF (140ml) and tetrabutylammonium iodide (7.92g) and sodium azide (3.065g) added.

20

The mixture was heated under N<sub>2</sub>, in the dark at 60-70°C for 1h. Upon cooling, diethyl ether and water were added and the organic layer washed successively with 0.1N HCl, water and brine, and dried over MgSO<sub>4</sub>. Concentration *in vacuo* gave an oil which was purified by column chromatography on silica gel (3/1 hexane/ethyl acetate). The fractions with R<sub>f</sub> = 0.42 (3/1 hexane/ethyl acetate) pooled and concentrated to yield 1-azido-4-(butoxycarbonylamino)but-2-yne as a yellow oil (3.049g).

25

Preparation of 1-Amino-4-(1-iminoethylamino)but-2-yne 2.5hydrochloride

To 1-azido-4-(butoxycarbonylamino)but-2-yne (0.294g) in dry THF (7ml) was added triphenyl phosphine (0.386g) and water (38ml) and the mixture stirred overnight at room temperature. After this time S-benzyl-2-thioacetimidate hydrochloride (0.237g) and ethanol (4ml) were added and the mixture stirred overnight at room temperature. The mixture was concentrated *in vacuo*, the residue partitioned between water and ethyl acetate, and the aqueous layer concentrated *in vacuo*. The residue was treated with ethanol and re-concentrated *in vacuo* (x2). The residue was triturated with diethyl ether and the ether decanted off (x2). The remaining solid was sucked dry on an oil pump to yield 1-butoxycarbonylamino-4-(1-iminoethylamino)but-2-yne hydrochloride as a white hygroscopic solid (0.332g).

1-Butoxycarbonylamino-4-(1-iminoethylamino)but-2-yne (0.332g) was deprotected by treatment with 4NHCl/dioxane (3ml) at 0°C and stirring at 0°C for 1h then at room temperature for 1h. After this time the mother liquor was decanted off and the sticky residue subjected to an oil pump vacuum to remove residual solvent. The resulting material was triturated with dry diethyl ether (x2), filtered off and washed with more dry diethyl ether. The yellow solid was finally dried in a drying pistol to yield 1-amino-4-(1-iminoethylamino)but-2-yne 2.5hydrochloride (0.184g), Mpt 182-185°C(dec), FAB M.S.  $M+1^+$  126.

Example 64

Preparation of 1-Amino-4-(1-imino-2-fluoroethylamino)but-2-yne dihydrobromide

1-Amino-4-(1-imino-2-fluoroethylamino)but-2-yne dihydrobromide was made from 1-azido-4-(butoxycarbonylamino)but-2-yne and S-benzyl-2-fluorothioacetimidate hydrobromide to yield 1-butoxycarbonylamino-4-(1-imino-2-fluoro-ethylamino)but-2-yne hydrobromide as an intermediate.

1-Butoxycarbonylamino-4-(1-imino-2-fluoroethylamino)but-2-yne hydrobromide (500mg) was deprotected by dissolving in glacial acetic acid (10ml), cooling until the solution almost solidified and adding HBr in acetic acid (45% w/v, 1.8ml). The mixture was stirred with cooling for 1h then at room temperature for 1h. After this time the mother liquor was decanted off and the sticky residue subjected to an oil pump vacuum to remove residual solvent. The resulting material was triturated with dry diethyl ether (x2), filtered off and washed with more dry diethyl ether. The yellow solid was recrystallised from ethanol/diethyl ether and finally dried in a drying pistol to yield 1-amino-4-(1-imino-2-fluoroethylamino)but-2-yne dihydrobromide, Mpt 174-176.5oC(dec.), FAB M.S. M+1<sup>+</sup> 144.

#### Example 65

##### Preparation of 1-(2-Aminoethoxy)-2-N-(1-iminoethyl)ethylamine 2hydrochloride

##### Preparation of N-(*tert*butoxycarbonyl)-2-(2-aminoethoxy)ethyl azide

The title compound was prepared from 2-(2-aminoethoxy)ethanol using the procedures described for equivalent non-ether linked straight chain aminoalcohols by Mattingly (*Synthesis*, 1990, 366-368).

##### Preparation of 1-(2-Aminoethoxy)-2-N-(1-iminoethyl)ethylamine 2hydrochloride

To N-(*tert*butoxycarbonyl)-2-(2-aminoethoxy)ethyl azide (0.315g) in dry THF (10ml) was added polymer supported triphenyl phosphine (0.46g) and water (37μl) and the mixture slowly stirred overnight at room temperature. After this time S-benzyl-2-thioacetimidate hydrochloride (0.352g) and ethanol (5ml) were added and the mixture stirred overnight at room temperature. The mixture was concentrated *in vacuo*, the residue partitioned between water and ether, and the aqueous layer concentrated *in vacuo* to furnish N-(*tert*butoxycarbonyl)-1-(2-Aminoethoxy)-2-N-(1-iminoethyl) ethylamine as a white foam.

Treatment of a solution of N-(*tert*butoxycarbonyl)-1-(2-Aminoethoxy)-2-N-(1-

iminoethyl) ethylamine (280mg) in dioxan with 10ml of 4N HCl in dioxan served to deprotect the molecule by stirring at room temperature for 1h. At this time a 50ml portion of ether and the suspension were vigorously stirred. The title compound was furnished as a hygroscopic solid on filtering this suspension and drying *in vacuo*.

FAB MS  $M+1^+$ , 100%,  $^1H$  NMR (360 MHz,  $D_2O$ )  $\delta_H$ , ppm 2.28 (1H, s, Me); 3.25 (2H, t,  $CH_2$ ), 3.55 (2H, t  $CH_2$ ), 3.82 (4H, m, 2x  $CH_2$ ), IR  $\nu_{max}$  (KBr disc) 3 111, 1 676, 1 643  $cm^{-1}$

## 10 Biological Data

NO synthase inhibition was determined by the following procedure:

### Purification of NOS from human placenta

15 Amion and chorion were removed from fresh placenta, which was then rinsed with 0.9% NaCl. The tissue was homogenised in a Waring blender in 3 volumes of HEDS buffer (20 mM Hepes pH 7.8, 0.1 mM EDTA, 5mM DTT, 0.2M sucrose) plus 0.1 mM PMSF. The homogenate was filtered through cheesecloth and then centrifuged at 1000g for 20 min. The supernatant was recentrifuged at 27,500g  
20 for 30 min. Solid ammonium sulphate was added to the supernatant to give 32% saturation. Precipitated protein was pelleted at 25,000g and then redissolved in a minimal volume of HEDS buffer plus 0.1 mM PMSF, 10 $\mu$ g/ml leupeptin and soybean trypsin inhibitor, and 1 $\mu$ g/ml pepstatin. The redissolved pellet was centrifuged at 15,000g for 10 min. To the supernatant was added 1/20 volume at  
25 2,5' ADP agarose resin (Sigma), and the slurry was mixed slowly overnight. In the morning, slurry was packed into a column. The resin was sequentially washed with HEDS, 0.5M NaCl in HEDS, HEDS, and then NOS was eluted with 10 mM NADPH in HEDS. The enzyme could be concentrated by ultra filtration and quick frozen and stored at -70°C without loss in activity for at least 3 months.

#### Assay for human placental NOS

NOS was assayed for the formation of citrulline following the procedure of Schmidt et al (PNAS 88 365-369, 1991) with these modifications: 20 mM Hepes, pH 7.4, 10 $\mu$ g/ml calmodulin, 2.5 mM CaCl<sub>2</sub>, 2.5 mM DTT, 125 $\mu$ M NADPH, 10 $\mu$ M tetrahydrobiopterin, 0.5mg/ml BSA, and 1 $\mu$ M L-[<sup>14</sup>C]arginine (New England Nuclear). Linearity of NOS-catalysed rate was confirmed prior to kinetic studies that used single time point determination of rate.

#### Purification of NOS from cytokine-induced human colorectal adenocarcinoma

##### DLD-1 cells

DLD-1 (ATCC No. CCL 221) were grown at 37°C, 5% CO<sub>2</sub> in RPMI 1640 medium supplemented with L- glutamine, penicillin, streptomycin, and 10% heat-inactivated fetal bovine serum. Cells were grown to confluence and then the following cocktail of cytokines were added: 100 units/ml interferon-gamma, 200 units/ml interleukin-6, 10 ng/ml tumour necrosis factor, and 0.5 ng/ml interleukin-1 $\beta$ . At 18-24 hr post-induction, cells were harvested by scraping and washed with phosphate-buffered saline. Pelleted cells were stored at -70°C. Purification of the induced NOS was performed at 4°C. Crude extract was prepared by three cycles of freeze/thawing cells in TDGB (20 mM tris pH 7.5, 10% glycerol, 1mM DTT, 2  $\mu$ M tetrahydrobiopterin). Extract was applied directly onto a column of 2',5' ADP sepharose (Pharmacia). Resin was sequentially washed with TDGB, 0.5M NaCl in TDGB, TDGB. NOS was eluted with 2 mM NADPH in TDGB. BSA was immediately added to give a final concentration of 1 mg/ml. NOS could be quick frozen and stored at -70°C without loss in activity for at least 2 months.

#### Assay for inducible human NOS

The formation of citrulline were assayed as described above except that 10 $\mu$ M FAD was included and calmodulin and CaCl<sub>2</sub> excluded from the assay mix.




Purification of NOS from human brain

Human brain NOS was prepared using variations of the procedures of Schmidt et al. (PNAS 88 365-369, 1991), Mayer et al. (Fed. Eur. Biochem. Soc. 288 187-191, 1991), and Bredt and Snyder, (PNAS 87 682-685, 1990). Briefly, fresh whole brains (3 with myelinated tissue dissected away, 1050g) were homogenised in cold buffer A (50 mM HEPES, pH 7.5 (pH at RT) and 0.5 mM EDTA, 10 mM DTT, 3.6 L total volume) with a polytron. The mixture was centrifuged at 13,000g for 1 hour and the supernatant fluid was removed (about 2050ml). To the supernatant fluid, solid ammonium sulphate (365g, about 30% of saturation) was added and stirred slowly for a total of 30 minutes. The precipitate was pelleted at 13,000g for 30 minutes and the pellet was resuspended in ~400ml of buffer A with 4 $\mu$ M tetrahydrobiopterin, 1 $\mu$ M FAD (Sigma), and 1 $\mu$ M FMN (Sigma). The solution was centrifuged at 41,000g for 60 minutes. The supernatant was removed, frozen by pouring into liquid nitrogen, and stored overnight at -70°C. The mixture was thawed and passed through at 2',5' ADP-agarose column (0.4g swelled in buffer A) at 4ml/min. The column was washed with 100ml buffer A, 200ml buffer A with 500 mM NaCl, 100ml Buffer A, then 30ml buffer A with 5 mM NADPH. To the enzyme solution was added tetrahydrobiopterin to 10 $\mu$ M, FAD and FMN to 1 $\mu$ M, and Tween to 0.1%. This solution was concentrated by Centriprep-30 to a volume of approximately 500 $\mu$ l. Enzyme activity was determined as described by Schmidt et al. 1991, except that 10 $\mu$ M tetrahydrobiopterin was included in the assay.

The compounds of the invention exhibit NOS inhibition properties, and certain groups of the compounds exhibit selective inhibition properties against either the inducible or neuronal NOS.

CLAIMS:

1. An acetamidine derivative of formula (I)




or a salt thereof, wherein

$\text{R}^1$  is hydrogen, a  $\text{C}_{1-6}$  hydrocarbyl group optionally substituted by halo, halo, nitro, cyano or a group  $\text{XR}^3$  wherein X is oxygen,  $\text{C}(\text{O})_m$  wherein m is 1 or 2,  $\text{S}(\text{O})_n$  wherein n is 0, 1 or 2, or a group  $\text{NR}^4$  wherein  $\text{R}^4$  is hydrogen or  $\text{C}_{1-6}$  alkyl; and  $\text{R}^3$  is hydrogen,  $\text{C}_{1-6}$  alkyl, or a group  $\text{NR}^5\text{R}^6$  wherein  $\text{R}^5$  and  $\text{R}^6$  are independently hydrogen or  $\text{C}_{1-6}$  alkyl, provided that  $\text{R}^3$  is not  $\text{NR}^5\text{R}^6$  when X is oxygen or  $\text{S}(\text{O})_n$ ;

$\text{R}^{1a}$  and  $\text{R}^{1b}$  are independently selected from hydrogen or halo;


$\text{R}^2$  is a  $\text{C}_{1-14}$  hydrocarbyl group which may optionally contain one or two heteroatoms, the group  $\text{R}^2$  being optionally substituted by one or more groups independently selected from halo;  $\text{N}_3$ ; nitro;  $\text{CF}_3$ ;  $\text{ZR}^7$  wherein Z is oxygen,  $\text{C}(\text{O})_{m'}$  wherein  $m'$  is 1 or 2,  $\text{S}(\text{O})_{n'}$  wherein  $n'$  is 0, 1 or 2, or a group  $\text{NR}^8$  wherein  $\text{R}^8$  is hydrogen or  $\text{C}_{1-6}$  alkyl and  $\text{R}^7$  is hydrogen,  $\text{C}_{1-6}$  alkyl or a group  $\text{NR}^9\text{R}^{10}$  wherein  $\text{R}^9$  and  $\text{R}^{10}$  are independently hydrogen or  $\text{C}_{1-6}$  alkyl; or  $\text{R}^2$  is substituted by a group



wherein  $R^{11}$  has a definition the same as for  $R^1$ ;

with the proviso that when  $R^1$  is a  $C_{1-6}$  alkyl group and  $R^2$  is a  $C_{1-14}$  hydrocarbyl substituted by two groups  $ZR^7$  wherein one group  $ZR^7$  is  $CO_2H$ , the other group  $ZR^7$  is not  $NH_2$ .

2. An acetamidine derivative as claimed in claim 1 wherein  $R^1$  is hydrogen, a  $C_{1-6}$  hydrocarbyl group optionally substituted by halo, halo, nitro, cyano or a group  $XR^3$  wherein  $X$  is oxygen,  $C(O)_m$  wherein  $m$  is 1 or 2,  $S(O)_n$  wherein  $n$  is 0, 1 or 2, or a group  $NR^4$  wherein  $R^4$  is hydrogen or  $C_{1-6}$  alkyl; and  $R^3$  is hydrogen,  $C_{1-6}$  alkyl, or a group  $NR^5R^6$  wherein  $R^5$  and  $R^6$  are independently hydrogen or  $C_{1-6}$  alkyl, provided that  $R^3$  is not  $NR^5R^6$  when  $X$  is oxygen or  $S(O)_n$ ; and  $R^{1a}$  and  $R^{1b}$  are independently selected from hydrogen or halo.
3. An acetamidine derivative as claimed in claim 1 or claim 2 wherein  $R^2$  is a  $C_{1-14}$  hydrocarbyl group which may optionally contain one or two heteroatoms, the group  $R^2$  being optionally substituted by one or two groups which may be the same or different, and are selected from halo;  $ZR^7$  wherein  $Z$  is oxygen,  $C(O)_m$  wherein  $m'$  is 1 or 2,  $S(O)_n$  wherein  $n'$  is 0, 1 or 2, or a group  $NR^8$  wherein  $R^8$  is hydrogen or  $C_{1-6}$  alkyl and  $R^7$  is hydrogen,  $C_{1-6}$  alkyl or a group  $NR^9R^{10}$  wherein  $R^9$  and  $R^{10}$  are independently hydrogen or  $C_{1-6}$  alkyl; or  $R^2$  is substituted by a group




wherein  $R^{11}$  has a definition the same as for  $R^1$ ;

with the proviso that when  $R^1$  is a  $C_{1-6}$  alkyl group and  $R^2$  is a  $C_{1-14}$  hydrocarbyl substituted by two groups  $ZR^7$  wherein one group  $ZR^7$  is  $CO_2H$ , the other group  $ZR^7$  is not  $NH_2$ .

5 4. An acetamidine derivative as claimed in any one of claims 1 to 3 wherein  $R^1$  is hydrogen,  $C_{1-4}$  alkyl, halo, nitro,  $NR^4R^3$  or  $SR^3$  wherein  $R^3$  and  $R^4$  are as defined in claim 1.

10 5. An acetamidine derivative as claimed in any one of claims 1 to 4 wherein  $R^2$  is a  $C_{1-8}$  hydrocarbyl group optionally containing one or two heteroatoms, each group  $R^2$  being optionally substituted by one group  $ZR^7$  wherein  $Z$  and  $R^7$  are as defined in claim 1, or a group




15

wherein  $R^{12}$  is fluoro, hydrogen or  $NH_2$ .


20 6. An acetamidine derivative as claimed in any one of claims 1 to 5 wherein  $R^1$  is hydrogen, halo or  $NH_2$ .

7. An acetamidine derivative as claimed in any one of claims 1 to 6 wherein  $R^2$  is a group:

25




or




wherein p, q and r are independently 0, 1 or 2 and R<sup>12</sup> is as defined in claim 1.

8. An acetamidine derivative as claimed in claim 1 of the formula (Ia):




10  
wherein R<sup>1</sup>, R<sup>1a</sup> and R<sup>1b</sup> are as hereinbefore defined, and R<sup>13</sup> is a C<sub>1-13</sub> hydrocarbyl group which may optionally contain one or two heteroatoms; the group R<sup>13</sup> being optionally substituted by one or two groups which may be the same or different, and are selected from halo, ZR<sup>7</sup> or a group




20  
wherein Z, R<sup>7</sup> and R<sup>11</sup> are as defined in claim 1.

9. A compound of formula (IB):




wherein R<sup>1</sup>, R<sup>1a</sup> and R<sup>1b</sup> are as hereinbefore defined, and R<sup>14</sup> represents one or two substituents on the phenyl ring independently selected from halo, ZR<sup>7</sup> or a group




wherein Z, R<sup>7</sup> and R<sup>11</sup> are as hereinbefore defined, or R<sup>14</sup> is a C<sub>1-8</sub> hydrocarbyl group optionally containing one or two heteroatoms, and optionally substituted by one or two groups which may be the same or different, and are selected from halo, ZR<sup>7</sup> or a group

5



10. A compound of the formula IC:

10



wherein R<sup>1</sup>, R<sup>1a</sup> and R<sup>1b</sup> are as defined in claim 1;

15


R<sup>15</sup> is a bond or CH<sub>2</sub>; and

Y is C<sub>1-6</sub> hydrocarbyl group, phenyl, indanyl or a 5-10 membered heterocyclic group containing 1 or 2 heteroatoms independently selected from O, N, and S;

20

wherein Y is optionally substituted by one or more groups independently selected from halo, amino, cyano, nitro, hydroxy, C<sub>1-6</sub> hydrocarbyl optionally substituted by hydroxy, C<sub>1-6</sub> alkoxy, C<sub>1-6</sub> thioalkyl, CF<sub>3</sub>, N<sub>3</sub>, a group:


25



wherein n is as defined in claim 1, C(O)<sub>m</sub>B or S(O)<sub>n</sub>B wherein m' and n' are as

defined in claim 1 and B is amino or C<sub>1-6</sub> alkyl, a group:

5




10 wherein n is as hereinbefore defined and D and E are independently hydrogen, hydroxy, amino, C<sub>1-6</sub> alkyl, -CNHCH<sub>3</sub>, -CO<sub>2</sub> tBu, or a 5-10 membered heterocyclic group containing 1 or 2 heteroatoms independently selected from O, N or S, a group:

15




wherein n and n' are as hereinbefore defined and F is phenyl, or a 5- or 6-membered heterocyclic group containing 1 or 2 heteroatoms selected from O, N and S optionally substituted by

20



25

a group:



wherein n is as hereinbefore defined and G is -CO<sub>2</sub>H, -CO<sub>2</sub>C<sub>1-6</sub> alkyl or a 5-10 membered heterocyclic group containing 1 or 2 heteroatoms independently selected from O, N or S optionally substituted by amino, CNHNH<sub>2</sub> and CH<sub>2</sub>SCNHNH<sub>2</sub>.

wherein Z, R<sup>7</sup> and R<sup>11</sup> are as defined in claim 1.


11. A compound as claimed in claim 10 wherein:

5 R<sup>15</sup> is a bond or CH<sub>2</sub>;

Y is C<sub>1-6</sub> hydrocarbyl, phenyl, indanyl or a 5-10 membered heterocyclic group containing 1 or 2 heteroatoms independently selected from O, N or S;


10 wherein Y is optionally substituted by one or more groups independently selected from halo, amino, cyano, nitro, hydroxy, C<sub>1-6</sub> hydrocarbyl optionally substituted by hydroxy, C<sub>1-6</sub> thioalkyl, a group:

15




wherein n is as defined in claim 1, C(O)<sub>m</sub>B or S(O)<sub>n</sub>B wherein m' and n' are as defined in claim 1 and B is amino or C<sub>1-6</sub> alkyl, a group:

20




25 wherein n is as hereinbefore defined and D and E are independently hydrogen, hydroxy, amino, C<sub>1-6</sub> alkyl, -CNHCH<sub>3</sub>, -CO<sub>2</sub>Bu, or a 5-10 membered heterocyclic group containing 1 or 2 heteroatoms independently selected from O, N or S, a group:





wherein n and n' are as hereinbefore defined and F is phenyl, or a 5- or 6-  
 5 membered heterocyclic group containing 1 or 2 heteroatoms selected from O, N and  
 S optionally substituted by



10

a group:



15

wherein n is as hereinbefore defined and G is a 5-10 membered heterocyclic group  
 containing 1 or 2 heteroatoms independently selected from O, N or S optionally  
 substituted by amino, CNH<sub>2</sub>NH<sub>2</sub> and CH<sub>2</sub>SCNHNH<sub>2</sub>.

20

12. A compound as claimed in claim 11 wherein:


R<sup>15</sup> is a bond or CH<sub>2</sub>:

Y is C<sub>1-6</sub> hydrocarbyl, phenyl, indanyl or a 5-10 membered heterocyclic group  
 containing 1 or 2 heteroatoms independently selected from O, N or S;

25


wherein Y is optionally substituted by one or more groups independently selected  
 from amino, C<sub>1-6</sub> hydrocarbyl optionally substituted by hydroxy, a group:

72



5

wherein n is as defined in claim 1, a group:



10


wherein n is as hereinbefore defined and D and E are independently hydrogen, hydroxy, amino, C<sub>1-6</sub> alkyl, -CNHCH<sub>3</sub>, or a 5-10 membered heterocyclic group containing 1 or 2 heteroatoms independently selected from O, N or S, a group:

15



wherein n and n' are as defined in claim 1 and F is phenyl, or a 5- or 6-membered heterocyclic group containing 1 or 2 heteroatoms selected from O, N and S optionally substituted by

20



25

a group:



wherein n is as hereinbefore defined and G is a 5-10 membered heterocyclic group

containing 1 or 2 heteroatoms independently selected from O, N or S, CNHNH<sub>2</sub> and CH<sub>2</sub>SCNHNH<sub>2</sub>.

5

13. A compound as claimed in claim 10 wherein R<sup>15</sup> is a bond.

14. A compound as claimed in claim 13 wherein Y is optionally substituted phenyl or an optionally substituted 5-10 membered heterocyclic group containing 1 or 2 heteroatoms independently selected from O, N or S.

10

15. A compound as claimed in claim 14 which is:

15

N-(3-(Hydroxymethyl)phenyl)Acetamidine;

N-(3-(Aminomethyl)phenyl)acetamidine;

N-(3-((1,2,3,4-tetrahydroisoquinol-2-yl)methyl)phenyl)-acetamidine;

N-(3-((Methylamino)methyl)phenyl)acetamidine;

N-(Dimethylamino)methyl)phenyl)acetamidine;

N-(3-(Aminomethyl)phenyl)-2-fluoroacetamidine;

N-(3-Morpholinomethyl)phenyl)acetamidine;

N-(3-(2-Aminoethyl)phenyl)acetamidine; or

20

N-(1,2,3,4-Tetrahydroisoquinolin-7-yl)acetamidine.

16. A compound as claimed in claim 10 wherein R<sup>15</sup> is -CH<sub>2</sub>-.

25

17. A compound as claimed in claim 16 wherein Y is optionally substituted phenyl, indanyl or a optionally substituted 5-10 membered heterocyclic group containing 1 or 2 heteroatoms independently selected from O, N or S.

18. A compound as claimed in claim 17 which is:

- N-(3-(aminomethyl)benzyl)acetamidine;  
N-(4-(aminomethyl)benzyl)acetamidine;  
N-(3-(Aminomethyl)benzyl)-2-fluoroacetamidine;  
N-(3-(Aminomethyl)benzyl)-2-aminoacetamidine;  
5 N-[[6-(Aminomethyl)-2-pyridyl]methyl]acetamidine;  
N-(3-(Aminomethyl)benzyl)-2,2,2-trifluoroacetamidine;  
N-((1H-Indazol-6-yl)methyl)acetamidine; or  
N-((3-Amino-5-indanyl)methyl)acetamidine.

- 10 19. A compound as claimed in claim 18 which is:

- N-(3-(aminomethyl)benzyl)acetamidine;  
N-(4-(aminomethyl)benzyl)acetamidine;  
N-(3-(Aminomethyl)benzyl)-2-fluoroacetamidine;  
15 N-(3-(Aminomethyl)benzyl)-2-aminoacetamidine;  
N-[[6-(Aminomethyl)-2-pyridyl]methyl]acetamidine;  
N-((1H-Indazol-6-yl)methyl)acetamidine; or  
N-((3-Amino-5-indanyl)methyl)acetamidine;

- 20 or a salt thereof.

20. A compound as claimed in claim 1 or 10 which is:

- N-(3-(aminomethyl)benzyl)acetamidine;  
25 N-(3-(aminomethyl)benzyl)-2-fluoroacetamidine;  
N-(4-(aminomethyl)benzyl)acetamidine;  
N-(4-aminobutyl)acetamidine;  
N-(5-aminopentyl)acetamidine;  
N-(6-aminohexyl)acetamidine;

- N-(7-aminoheptyl)acetamidine;  
N,N'-(1,3-phenylenebis (methylene))bis(acetamidine);  
N,N'-(1,4-phenylenebis (methylene))bis(acetamidine);  
N,N'-(1,3-phenylene (diethylene))bis(acetamidine);  
5 N,N'-(1,4-phenylene (diethylene))bis(acetamidine);  
N-(3-(aminomethyl)benzyl)-2-methoxyacetamidine;  
2-Amino-N-(3-(aminomethyl)benzyl)acetamidine;  
N-(3-(hydroxymethyl)phenyl)acetamidine;  
N-(3-(aminomethyl)phenyl)acetamidine;  
10 N-(3-(Aminomethyl)phenyl)acetamidine;  
N-(3-(Aminomethyl)phenyl)-2-fluoroacetamidine; or  
N-((3-Amino-5-indanyl)methyl)acetamidine;

or a salt thereof.

15

21. A compound as claimed in claim 20 which is:

- N-(3-(aminomethyl)benzyl)-2-fluoroacetamidine;  
N-(3-(Aminomethyl)phenyl)acetamidine;  
20 N-(3-(Aminomethyl)phenyl)-2-fluoroacetamidine; or  
N-((3-Amino-5-indanyl)methyl)acetamidine;

or a salt thereof.

25

22. A compound as defined in any one of claims 1 to 21 for use in medicine.

23. A pharmaceutical formulation comprising at least one compound as defined in any one of claims 1 to 21 together with one or more pharmaceutically acceptable carriers, diluents or excipients.

24. The use of a compound as defined in any one of claims 1 to 21 in the manufacture of a medicament for the prophylaxis or treatment of a condition requiring inhibition of NO synthase.
- 5 25. The use as claimed in claim 24 wherein the compound inhibits one of the NO synthase isoenzymes with little or no inhibition of the other isoenzymes.
26. A method of treating a condition requiring inhibition of NO synthase, which comprises the step of administering to a mammal in need thereof an effective  
10 amount of a compound as defined in any one of claims 1 to 21 or a pharmaceutical formulation as claimed in claim 23.
27. A method as claimed in claim 26 wherein the compound, or pharmaceutical formulation, inhibits one NO synthase isoenzyme with little or no inhibition of  
15 other isoenzymes.
28. A method as claimed in claim 26 or claim 27 wherein the mammal is a human.
- 20 29. The use of a compound as defined in any one of claims 1 to 21 in the manufacture of a medicament for the prophylaxis or treatment of shock states resulting from overproduction of NO by iNOS.
- 25 30. The use as claimed in claim 29 wherein the condition is septic shock, shock caused by fulminant hepatic failure, shock caused by therapy with one or more cytokines or shock caused by therapy with cytokine-inducing agents.
31. The use as claimed in claim 30 wherein the condition is septic shock.

32. The use of a compound as defined in any one of claims 1 to 21 in the manufacture of a medicament for the prophylaxis or treatment of an auto-immune disease or an inflammatory disease.
- 5 33. The use as claimed in claim 32 wherein the disease is rheumatoid arthritis, osteoarthritis, ulcerative colitis, gastritis, mucosal inflammation resulting from infection, adult respiratory distress syndrome, asthma, myocarditis, multiple sclerosis, diabetes melitus, glomerulonephritis, dermatitis, psoriasis, urticaria, transplant organ rejection or systemic lupus erythematosus.
- 10 34. The use as claimed in any one of claims 29 to 33 wherein the compound is as defined in claim 18 or claim 19.
- 15 35. The use of a compound as defined in any one of claims 1 to 21 in the manufacture of a medicament for the prophylaxis or treatment of a condition resulting from the overproduction of NO by nNOS.
- 20 36. The use as claimed in claim 35 wherein the condition is cerebral ischemia, CNS trauma, epilepsy, AIDS dementia, chronic neurodegenerative disease, chronic pain, priapism, obesity or hyperphagia.
37. The use as claimed in claim 36 wherein the compound is as defined in claim 15.
- 25 38. The use of a compound as defined in any one of claims 1 to 21 in the manufacture of a medicament for the prophylaxis or treatment of a condition which requires inhibition of both iNOS and nNOS.
39. A method for the prophylaxis or treatment of a condition requiring inhibition

of iNOS which comprises the step of administering to a mammal an effective amount of a compound as defined in any one of claims 1 to 21.

5 40. A method as claimed in claim 39 wherein the condition is septic shock, shock caused by fulminant hepatic failure, shock caused by therapy with one or more cytokines or shock caused by therapy with cytokine-inducing agents.

41. A method as claimed in claim 40 wherein the condition is septic shock.

10 42. A method as claimed in claim 39 wherein the condition is an auto-immune disease or an inflammatory disease.

15 43. A method as claimed in claim 42 wherein the condition is rheumatoid arthritis, osteoarthritis, ulcerative colitis, gastritis, mucosal inflammation resulting from infection, adult respiratory distress syndrome, asthma, myocarditis, multiple sclerosis, diabetes melitus, glomerulonephritis, dermatitis, psoriasis, urticaria, transplant organ rejection or systemic lupus erythematosus.

20 44. A method as claimed in any one of claims 39 to 43 wherein the compound is as defined in claim 18 or claim 19.


25 45. A method for the prophylaxis or treatment of a condition requiring inhibition of nNOS which comprises the step of administering to a mammal an effective amount of a compound as defined in any one of claims 1 to 21.

46. A method as claimed in claim 45 wherein the condition is cerebral ischemia, CNS trauma, epilepsy, AIDS dementia, chronic neurodegenerative disease, chronic pain, priapism, obesity or hyperphagia.



47. A method as claimed in claim 45 or claim 46 wherein the compound is as defined in claim 15.


48. A process for the preparation of a compound of formula (I), as defined in claim 1 or claim 2 comprising the reaction of an amine of formula (II):



10

or a protected derivative thereof wherein R<sup>2</sup> is as hereinbefore defined, with a 1-benzylthioethaniminium derivative of formula (III):

15



wherein R<sup>1</sup> is as hereinbefore defined followed by deprotection if necessary.

# INTERNATIONAL SEARCH REPORT

Intern:      Application No  
PCT/GB 95/02978

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6    C07C257/14    C07D213/75    C07D333/20    C07C311/46    C07D217/18  
           C07D277/40    C07D217/04    C07D231/56    C07D231/12    C07D295/135  
           C07D521/00    A61K31/155

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6    C07C

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	CHEMICAL ABSTRACTS, vol. 102, no. 10, 11 March 1985 Columbus, Ohio, US; abstract no. 84808, J. OSZCZAPOWICZ ET AL. see RN 94793-54-1, Ethanimidamide, N-(3-bromophenyl)- see RN 82635-74-3, Ethanimidamide, N-(3-fluorophenyl)- see RN 35372-37-3, Ethanimidamide, N-(3-ethoxyphenyl)- see RN 35372-31-7, Ethanimidamide, N-(3-nitrophenyl)- & J. CHROMATOGR., vol. 315, 1984 pages 95-100,  --- -/--	1-6, 9-11, 13, 14

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

4 March 1996

Date of mailing of the international search report

08.03.96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+ 31-70) 340-3016

Authorized officer

Seufert. G

# INTERNATIONAL SEARCH REPORT

Intern al Application No  
PCT/GB 95/02978

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	JOURNAL OF MEDICINAL CHEMISTRY, vol. 21, no. 12, 1978 WASHINGTON US, pages 1283-90, W. S. SAARI ET AL. 'Trichloroacetamidines, a new class of positive inotropic agents' see table I ---	1-6,8, 10-12, 16,17
X	DE,A,26 01 137 (MERCK & CO INC) 22 July 1976  see claims; examples ---	1-6,8, 10,11, 16,17
X	DE,A,23 00 221 (AGFA GEVAERT AG) 18 July 1974  see page 3, compound II and examples ---	1-3,5, 9-11,13, 14
X	CHEMICAL ABSTRACTS, vol. 68, no. 21, 20 May 1968 Columbus, Ohio, US; abstract no. 95274, T. T. CONWAY ET AL. see abstract see RN 20211-50-1, Ethanimidamide, 2-mercapto-N-(2- pyridinylmethyl)-, dihydrochloride see RN 19412-58-9, Acetamidine, 2-mercapto-N-[2-(2- pyridyl)ethyl]-, dihydrochloride see RN 19412-57-8, Acetamidine, 2-mercapto-N-[(6-meth yl-2-pyridyl)methyl]-. dihydrochloride & J. PHARM. SCI., vol. 57, no. 3, 1968 pages 455-9, ---	1-4,8, 10-12, 16,17
X	JOURNAL OF HETEROCYCLIC CHEMISTRY, vol. 3, no. 4, 1966 PROVO US, pages 472-5, A. P. PARULKAR ET AL. 'N-Heteroaralkyl substituted .alpha.-amidinium thiolsulfates' see page 474, left column, line 5 - right column, line 10 ---	1-6,8, 10-12, 16,17
X	JOURNAL OF MEDICINAL CHEMISTRY, vol. 22, no. 3, 1979 WASHINGTON US, pages 295-301, W. A. BOLHOFF ET AL. '2-Mercaptoacetamidines as gastric antisecretory agents' see tables I, II, III ---	1-6, 8-14,16, 17

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 95/02978

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	JOURNAL OF THE CHEMICAL SOCIETY, PERKIN TRANSACTIONS 1, 1976 LETCHWORTH GB, pages 1847-52, W. F. KEIR ET AL. 'Amidinoacetamides in the synthesis of pyrimidines, imidazoles and purines' see page 1848, compounds 1b-d ---	1-3,5,8, 10-12, 16,17
X	DE,A,20 20 155 (ISHIHARA SANGYO KAISHA) 4 November 1971  see page 6 - page 9 ---	1-6, 9-11,13, 14
X	DE,A,25 57 651 (BASF AG) 23 June 1977  see claim 2; examples ---	1-6, 9-11,13
X	GB,A,1 183 135 (CIBA) 4 March 1970  see page 9, left column, line 17 - line 18 ---	1-8, 10-12, 16,17
X	DATABASE CROSSFIRE Beilstein Informationssysteme GmbH, Frankfurt DE see BRN 2346409 & CHEM. BER., vol. 93, 1960 pages 2230-38, ---	1-5, 10-13,16
X	DATABASE CROSSFIRE Beilstein Informationssysteme GmbH, Frankfurt DE see BRN 3777937 & JUSTUS LIEBIGS ANN. CHEM., vol. 575, 1952 page 90ff ---	1-6,8, 10-13,16
X	DATABASE CROSSFIRE Beilstein Informationssysteme GmbH, Frankfurt DE see BRN 7088281 see BRN 7089080 & FARMACO ED. SCI., vol. 43, no. 1, 1988 pages 103-112, ---	1-3,5,8, 10-13,16
X	DATABASE CROSSFIRE Beilstein Informationssysteme GmbH, Frankfurt DE see BRN 1790943 & J. CHEM. SOC., 1947 page 730 ---	1-3,5,8, 10-13,16

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/GB 95/02978

C(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE CROSSFIRE Beilstein Informationssysteme GmbH, Frankfurt DE see BRN 394415 &amp; CAN. J. CHEM., vol. 39, 1961 pages 761-764,</p> <p style="text-align: center;">---</p>	1-4,6, 10-14
X	<p>DATABASE CROSSFIRE Beilstein Informationssysteme GmbH, Frankfurt DE see BRN 3278146 &amp; RECL. TRAV. CHIM. PAYS-BAS, vol. 70, 1951 page 638,641</p> <p style="text-align: center;">---</p>	1-4,6, 9-11,13
X	<p>DATABASE CROSSFIRE Beilstein Informationssysteme GmbH, Frankfurt DE see BRN 2717900 &amp; GB,A,861 526 1958</p> <p style="text-align: center;">---</p>	1-6, 9-11,13
X	<p>DATABASE CROSSFIRE Beilstein Informationssysteme GmbH, Frankfurt DE see BRN 746863 &amp; BULL CHEM SOC. JPN, vol. 45, 1972 pages 2010-15,</p> <p style="text-align: center;">---</p>	1-6,8, 10-12, 16,17
X	<p>DATABASE CROSSFIRE Beilstein Informationssysteme GmbH, Frankfurt DE see BRN 473624 &amp; J. INDIAN CHEM. SOC., vol. 39, 1962 pages 208-210,</p> <p style="text-align: center;">---</p>	1-6, 10-14
X	<p>DATABASE CROSSFIRE Beilstein Informationssysteme GmbH, Frankfurt DE see BRN 113878 &amp; J. HETEROCYCL. CHEM., vol. 12, 1975 page 407</p> <p style="text-align: center;">---</p>	1-6, 10-14
X	<p>DATABASE CROSSFIRE Beilstein Informationssysteme GmbH, Frankfurt DE see BRN 6423899 &amp; GAZZ. CHIM.ITAL., vol. 113, no. 1-2, 1983 pages 77-82,</p> <p style="text-align: center;">---</p>	1-3,6, 10-13

## INTERNATIONAL SEARCH REPORT

Internat'l Application No

PC1/GB 95/02978

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
DE-A-2601137	22-07-76	AR-A- 213168	29-12-78
		AU-B- 499649	26-04-79
		AU-B- 1006076	14-07-77
		BE-A- 837569	14-07-76
		BG-A- 25074	12-07-78
		FR-A- 2297615	13-08-76
		GB-A- 1490671	02-11-77
		JP-A- 51125202	01-11-76
		LU-A- 74170	31-12-76
		NL-A- 7515256	19-07-76
		SE-A- 7514776	16-07-76
DE-A-2300221	18-07-74	BE-A- 809222	28-06-74
		FR-A,B 2322862	01-04-77
		GB-A- 1404442	28-08-75
		JP-A- 49095969	11-09-74
		US-A- 3931221	06-01-76
DE-A-2020155	04-11-71	NONE	
DE-A-2557651	23-06-77	NONE	
GB-A-1183135	04-03-70	BE-A- 713698	14-10-68
		CH-A- 511863	31-08-71
		CH-A- 523264	31-05-72
		CH-A- 527828	15-09-72
		DE-A- 1770135	23-09-71
		FR-M- 7495	08-12-69
		FR-A- 1588463	17-04-70
		FR-M- 7496	08-12-69
WO-A-9313055	08-07-93	AU-B- 3169293	28-07-93
		EP-A- 0618898	12-10-94
		HU-A- 70502	30-10-95
		JP-T- 7502512	16-03-95
		ZA-A- 9210018	23-06-94

FURTHER INFORMATION CONTINUED FROM PCT/ISA/

Although claims 26 - 28, 39 - 47 are directed to a method of treatment of the human body the search has been carried out and based on the alleged effects of the compounds/compositions.

Claims searched completely: 15, 18 - 21, 34, 37, 44, 47

Claims searched incompletely: 1 - 14, 16 - 17, 22 - 33, 35 - 36, 38 - 43, 45 - 46, 48

The vast number of theoretically conceivable compounds resulting from the combination of all claimed substituents precludes a comprehensive search of all the claims. For economical reasons search and search report had to be limited. The documents cited in the report are just a minute and arbitray selection of relevant documents and/or compounds. The search report can only be regarded as complete for the explicitly disclosed compounds.

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/GB95/02978

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:  
  
Please see attached sheet ./.
2. ☐ Claims Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:  
  
Please see attached sheet ./.
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.



# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/GB 95/02978

## C(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO,A,93 13055 (WELLCOME FOUND) 8 July 1993 see claims	1-48
A	--- JOURNAL OF MEDICINAL CHEMISTRY, vol. 37, no. 23, November 1994 WASHINGTON US, pages 3886-88, -----	1-48

